



ISPA | Instituto Superior de Psicologia Aplicada

Adult neurogenesis in a new model specie, the  
cichlid fish *Oreochromis mossambicus*.

Magda Teles

Orientador de dissertação:

Professor Rui Oliveira

Coordenador de seminário de dissertação:

Professor Rui Oliveira

Tese submetida como requisito parcial para a obtenção do grau de:

Mestre em Psicobiologia

2010

Dissertação de Mestrado realizada sob a orientação de professor Rui Oliveira, apresentada no Instituto Superior de Psicologia Aplicada para obtenção de grau de Mestre na especialidade de Psicobiologia conforme despacho da DGES nº 6037 / 2007 publicado em Diário da Republica 2ª série de 23 de Março, 2007

## **Acknowledgments**

I would like to thank to my supervisor, Professor Rui Oliveira. He made this thesis possible giving me access to all the resources, and the most important, the scientific input that I needed to make this work real and above all, for keeping me motivated even when I'm not.

I would also like to thank to Professor Gunther Zupanc from Jacobs University, Bremen, for having me during four months in his lab. He taught me all the techniques used in the present work.

To David Gonçalves for his corrections and for everything that he taught me during the last years, I have learned a lot... Can't thank you enough!!

To Ushi for all the support that she gave me in Bremen, not only in the Lab!!!

To Ruxandra for the help with the confocal microscopy.

To Leonor, she helped me to get inspiration in moment where inspiration was in fairway kingdom!!!

To Marta she is always there with a positive and funny thing to say. She always believes!!

To Miguel for the reviewing that he did in the state of the art section. Now he knows everything that he always wanted to know about adult neurogenesis and he never had the courage to ask ...

To all my colleagues, they are always available for one more presentation and for the daily support!!!

To my family, friends and David for all the patience, whether I be in a good mood, tired, angry, or stressed... they are always there for me !!!!!!



## Resumo

Em comparação com outros vertebrados, os peixes teleósteos têm um enorme potencial para produzir células novas no cérebro de animais adultos. Em contraste com os mamíferos, onde o processo de neurogênese adulta encontra-se restrito a duas áreas cerebrais, a zona subventricular (SVZ) e a zona subgranular parte do giro dentado do hipocampo, em peixes teleósteos foram descritas mais de 10 regiões neurogênicas. Através da marcação de células mitóticas com 5-bromo-2'-deoxiuridina (BrdU), foram caracterizadas as zonas proliferativas da Tilapia de Moçambique (*Oreochromis mossambicus*). Nesta espécie, foram encontradas zonas proliferativas em regiões específicas do bolbos olfactivo, telencéfalo, região pré-óptica, hipotálamo, tálamo, tecto óptico, *torus longitudinalis*, nas três divisões do cerebelo, *valvula cerebelli*, *corpus cerebelli*, e *lobus caudalis* e na região da medula, abrangendo assim toda a extensão cerebral. A localização destas zonas proliferativas parece ser extremamente conservada ao longo da taxonomia e até o número total de células produzidas parece ser mantido com pouca variação. Com um tempo de sobrevivência de 2 horas, foram encontrados na tilapia um total de 80.000 células novas em comparação com as 100.000 descritas para o peixe eléctrico *Apteronotus leptorhynchus*. Os nossos resultados sugerem que a actividade mitótica em regiões discretas do cérebro adulto são uma característica primitiva que tem sido conservada ao longo da evolução.

Palavras Chave: Tilápia de Moçambique, Neurogênese Adulta, Bolbos olfactivos, Telencéfalo, Tecto óptico, Cerebelo, BrdU, Peixes teleósteos

## Abstract

Compared to other vertebrate species, fish exhibit an enormous potential to produce new cells in the adult brain. In contrast to mammals, where proliferation zones are restricted to two brain areas, the sub ventricular zone (SVZ), and the subgranular zone (SGZ), part of the dentate gyrus of hippocampus, in teleost species more than 10 neurogenic regions have been described. By labeling mitotically dividing cells with 5-bromo-2'-deoxyuridine (BrdU), we have characterized the proliferation zones in the Mozambique tilapia (*Oreochromis mossambicus*). Proliferation zones were located in specific brain regions of the olfactory bulb, telencephalon, preoptic area, hypothalamus, thalamus, optic tectum, torus longitudinalis, in all three subdivisions of the cerebellum, the valvula cerebelli, the corpus cerebelli, and the lobus caudalis cerebelli and in the region of the medulla oblongata. These proliferation zones appeared to be extremely conserved across taxonomy and even the total number of new generated cells seems to be preserved. After 2 hours survival time we found a total of approximately 80.000 new cells for tilapia compared to 100.000 new cells described for *Apteronotus leptorhynchus*. Our results suggest that the presence of mitotic activity in specific brain regions is a primitive feature that has been conserved through evolution.

**Key words:** Mozambique tilapia, adult neurogenesis, olfactory bulb, telencephalon, optic tectum, cerebellum, BrdU, teleost fish.

## Table of Contents

Acknowledgments.....	ii
Resumo.....	iv
Abstract .....	v
List of Figures .....	viii
List of Tables.....	x
Abbreviation list.....	xi
Introduction .....	1
Materials and methods .....	2
Animals .....	2
Intraperitoneal injection of 5-bromo-2'-deoxyuridine.....	3
Intracardial perfusion and tissue preparation .....	3
BrdU immunohistochemistry .....	4
Microscopic examination and data analysis.....	4
Results.....	5
Proliferation zones .....	5
Olfactory bulb .....	5
Telencephalon .....	10
Diencephalon.....	13
Optic tectum .....	14
Nucleus recessi lateralis .....	15
Cerebellum .....	16
Valvula cerebelli .....	16

Lateral part of the valvula cerebelli .....	16
Medial part of the valvula cerebelli .....	17
Corpus cerebelli .....	17
Lobus caudalis.....	18
Discussion .....	20
Cell proliferation .....	21
Telencephalon .....	21
Olfactory bulb .....	21
Dorsal and ventral telencephalon .....	23
Optic tectum .....	24
Cerebellum .....	26
Conclusion .....	27
Bibliography.....	28
Annex I.....	<b>Erro! Marcador não definido.</b>
State of the art .....	32
The development of the nervous system.....	32
The importance of the cell cycle .....	32
Adult neurogenesis.....	34
Adult neurogenesis a historical view .....	34
Adult neurogenesis in the mammalian brain.....	35
Adult neurogenesis in non-mammalian vertebrates .....	37
Bibliography.....	42

## List of Figures

Fig 1 - Distribution and semi-quantitative analysis of proliferation zones in the forebrain (a-c) and midbrain (d-f) of Mozambique tilapia. Mitotic zones were revealed by the analysis of BrdU labeling in coronal sections at a post-administration survival time of 2 hours. Proliferative areas are indicated by black dots. Each dot represents approximately 50 BrdU-labeled cells.

Fig 2 - Distribution and semi-quantitative analysis of proliferation zones in the hindbrain of Mozambique tilapia. Mitotic zones were revealed by the analysis of BrdU labeling in coronal sections at a post-administration survival time of 2 hours. Proliferative areas are indicated by black dots. Each dot represents approximately 50 BrdU-labeled cells.

Fig 3 - Confocal images of BrdU-labeled cells in coronal sections of the olfactory bulb and telencephalon in adult tilapia at 2 hours post-administration survival time. a: Olfactory bulb (BO). Relatively few labeled cells were found in this brain region. Most of these cells were located in the glomerular layer (BO<sub>gl</sub>), whereas only a few were observed in the granular layer (BO<sub>gra</sub>). The labeled cells in BO<sub>gl</sub> tended to occupy ventral and lateral regions (arrowhead). b: Anterior subdivision of the dorsal telencephalon (DMa). This brain region was distinguished by a high density of BrdU-labeled cells in a narrow zone at the ventricle (V). c: Medial part of the ventral subdivision of the ventral telencephalon (VVm). The majority of labeled cells were concentrated at the rostral pole of this structure, from which the section was taken. DLa, anterior subdivision of the dorsolateral telencephalon. d: Dorsal part of the ventral telencephalon (Vd). The labeled cells in this brain area are exclusively found in the immediate vicinity of the ventricle. DMvv, ventral subdivision of the ventral part of the dorsomedial telencephalon.

Fig 4 - Confocal images of BrdU-labeled cells in coronal sections of the diencephalon in adult tilapia at 2 hours post-administration survival time. a: Anterior part of the periventricular preoptic nucleus (PPa). BrdU labeled cells were found over the entire dorsoventral extent of the nucleus.

b: Periventricular nucleus of the posterior tuberculum (TPP). Throughout this area, labeled cells displayed a remarkably high packing density. CP, central posterior thalamic nucleus.

Fig 5 - Confocal images of BrdU-labeled cells in coronal sections of the optic tectum (TeO) in adult tilapia at 2 hours post-administration survival time. a: Rostral level of TeO. Most of the new cells were concentrated in the innermost layer (Layer 1). This layer intrudes the dorsolateral aspect of the torus longitudinalis (TL). The site of the intrusion is distinguished by the presence of a dense cluster of labeled cells (arrowhead). b: Caudal level of TeO. The vast majority of the new cells were concentrated, at a uniform and high areal density, in Layer 1.

Fig 6 - Confocal images of BrdU-labeled cells in coronal sections of the recessus lateralis (RL) in adult tilapia at 2 hours post-administration survival time. a: Lateral part of nucleus recessi lateralis (nRLl). A high areal density of BrdU-labeled cells was found at the ventral and ventrolateral aspects of this structure. b: Medial part of the nucleus recessi lateralis (nRLm). Labeled cells were equally distributed throughout the medial border of this structure. ILvm, ventromedial subdivision of the inferior lobe of the hypothalamus.

Fig 7- Confocal images of BrdU-labeled cells in coronal sections of the cerebellum in adult tilapia at 2 hours post-administration survival time. a: Lateral part of the valvula cerebelli (Val). The majority of the new generated cells were located in the molecular layer of Val (Val<sub>mol</sub>). By contrast, in the granular layer of this structure (Val<sub>gra</sub>) very few cells were observed. The distribution of the cells in Val<sub>mol</sub> was restricted to the dorsal and medial aspect of this structure, with the cells being mostly round and ovoid. Some of these cells, indicated by arrow, are shown at higher magnification in a'. At the lateral aspect of Val<sub>mol</sub>, a stripe of elongated cells was found along the dorsoventral extent. Some of these cells, indicated by arrowhead, are shown at higher magnification in a". b: Medial part of the valvula cerebelli (Vam). Virtually all of the BrdU-labeled cells were located in the molecular layer of Vam (Vam<sub>mol</sub>). Their nuclei displayed an elongated morphology and were orientated parallel to the midline. Part of these cells are shown at higher magnification in b'. c: Corpus cerebelli (CCe). The majority of labeled cells were found in the molecular layer of this structure (CCe<sub>mol</sub>) immediately beneath the pial surface. They

stretched over several hundreds of micrometers laterally in each hemisphere. In addition, BrdU-labeled cells occurred in the midline region of the dorsal molecular layer and extended from the dorsal pial surface to the dorsal aspect of the granular layer of the CCe (CCe<sub>gra</sub>). Part of the latter cellular population, indicated by arrow, is shown in c'.

## **List of Tables**

Table 1- Quantitative analysis of the proliferation zones in all brain regions of tilapia. Proliferative zones were revealed by the analysis of BrdU labeling at 2 hours survival time.

## Abbreviation list

A	anterior thalamic nucleus
AC	anterior commissure
BO	olfactory bulb
BO <sub>gl</sub>	glomerular layer of the olfactory bulb
BO <sub>gra</sub>	granular layer of the olfactory bulb
CC	crista cerebellaris
CCe	corpus cerebelli
CCe <sub>gra</sub>	granular layer of corpus cerebelli
CCe <sub>mol</sub>	molecular layer of corpus cerebelli
CE	central nucleus of the inferior lobe
Ch	horizontal commissure
CM	corpus mamillare
CP	central posterior thalamic nucleus
CVEn	ventral rhombencephalic commissure
D	dorsal telencephalon
DA	anterior part of the dorsal telencephalon
DC	dorsocentral telencephalon
DCa	anterior subdivision of the dorsal central telencephalon
DCad	dorsal part of the anterior subdivision of the dorsocentral telencephalon
DCm	medial subdivision of the dorsocentral telencephalon
DD	dorsal division of the dorsal telencephalon
DFl	nucleus diffusus lateralis of the inferior lobe
DFld	dorsal subdivision of nucleus diffusus lateralis of the inferior lobe
DFlv	ventral subdivision nucleus diffusus lateralis of the inferior lobe
DFm	nucleus diffusus medialis of the inferior lobe
DL	dorsolateral telencephalon
DLa	anterior subdivision of the dorsolateral telencephalon
DLd	dorsal subdivision of the dorsolateral telencephalon
DLp	posterior subdivision of the dorsolateral telencephalon
DLv	ventral subdivision of the dorsolateral telencephalon
DM	dorsomedial telencephalon
DMA	anterior subdivision of the dorsomedial telencephalon
DMdd	dorsal part of the dorsal subdivision of the dorsomedial telencephalon
DMdv	ventral part of the dorsal subdivision of the dorsomedial telencephalon
DMv	ventral subdivision of the dorsomedial telencephalon
DMvd	dorsal part of the ventral subdivision of the dorsomedial telencephalon
DMvv	ventral subdivision of the ventral part of the dorsomedial telencephalon

DP	dorsal posterior thalamic nucleus
Dp	posterior part of dorsal telencephalon
EG	eminentia granularis
G	glomerular nucleus
GC	central gray
H	habenula
HV	ventral zone of the periventricular hypothalamus
IL	inferior lobe of the hypothalamus
ILdl	dorsolateral subdivision of the hypothalamus
ILvm	ventromedial subdivision of the inferior lobe of the hypothalamus
IPn	interpeduncular nucleus
LCe	lobus caudalis
LCe <sub>gra</sub>	granular layer lobus caudalis
LH	lateral hypothalamic nucleus
LVII	lobus facialis
LX	lobus vagi
M	
MLF	medial longitudinal fascicle
mol	molecular layer
nIII	nucleus oculomotorius
nRL	nucleus recessi lateralis
nRLl	lateral part of nucleus recessi lateralis
nRLm	medial part of the nucleus recessi lateralis
nRP	nucleus recessi posterioris
nVd	descending trigeminal nucleus
nX	
P	preoptic region
PGm	medial part of the preglomerular nucleus
PP	periventricular preoptic nucleus
PPa	anterior part of the periventricular preoptic nucleus
PPm	medial part of the periventricular preoptic nucleus
PPp	posterior part of the periventricular preoptic nucleus
RL	recessus lateralis
RV	rhombencephalic ventricle
S	
Sc	suprachiasmatic nucleus
TA	nucleus anterior tuberis
TeO	optic tectum
TGN	tertiary gustatory nucleus
TGNI	lateral part of the tertiary gustatory nucleus

TL	torus longitudinalis
TLA	torus lateralis
TLAi	inferior subdivision of the torus lateralis
tO	optic tract
tOd	tractus opticus dorsalis
tolfm	medial part of the olfactory tract
tOv	tractus opticus ventralis
TP	nucleus posterior tuberculum
TPP	periventricular nucleus of the posterior tuberculum
TS	torus semicircularis
tTB	tectobulbar tract
V	ventral telencephalon
V	ventricle
Val	lateral part of valvula cerebelli
Val <sub>gra</sub>	granular layer of the lateral part of valvula cerebelli
Val <sub>mol</sub>	molecular layer of the lateral part of the valvula cerebelli
Vam	medial part of valvula cerebelli
Vam <sub>gra</sub>	granular layer of the medial part of the valvula cerebelli
Vam <sub>mol</sub>	molecular layer of the medial part of the valvula cerebelli
Vd	dorsal part of the ventral telencephalon
VM	ventromedial thalamic nucleus
Vs	supracommissural part of the ventral telencephalon
VVm	medial part of the ventral subdivision of the ventral telencephalon
XY	
Z	

## **Introduction**

It is now well established that neurogenesis occurs throughout adulthood in the mammalian brain (Gage, 2000). In mammals this process occurs only in two brain areas located at the telencephalon, the sub ventricular zone (SVZ) lining the lateral ventricles, from where the newly generated cells migrate along the so-called rostral migratory stream into the olfactory bulb (Altman, 1969; Alvarez-buylla, Ling, & Yu, 1994; Alvarezbuylla, Ling, & Yu, 1994) and the subgranular zone (SGZ), part of the dentate gyrus of hippocampus. Since this discovery, a lot of research has been directed towards understanding the physiology and the molecular mechanisms underlying neuronal differentiation. However, neurogenesis within the mature brain is not restricted to mammals, and affects other vertebrates in a large extent, including birds, reptiles, amphibians and fishes (Chapouton, Jagasia, & Bally-Cuif, 2007).

Like in the adult mammalian brain, in non-mammalian brains the majority of the mitotic cells are found in discrete areas called “proliferation zones” and most of these zones are located near the surfaces of ventricles or related systems. In birds, constitutive adult proliferation and neurogenesis are found dispersed along the telencephalic lateral ventricle, particularly in the ventral and dorsal aspects of the lateral wall of this structure (Alvarez-buylla et al., 1994; Alvarez-buylla, Theelen, & Nottebohm, 1990; Goldman & Nottebohm, 1983). In reptiles, post-natal proliferation have been found in several brain areas such as the olfactory bulb, striatum, dorsoventricular ridge, cortex, nucleus sphericus and cerebellum (Font, Desfilis, Perez-Canellas, & Garcia-Verdugo, 2001; García-Verdugo, Llahi, Ferrer, & Lopezgarcia, 1989; López-García, Molowny, Garciaverdugo, & Ferrer, 1988; Marchioro et al., 2005; Pérez-Sánchez, Molowny, Garciaverdugo, & Lopezgarcia, 1989). In amphibian brain, there is only one study that describes the proliferative spots (Raucci et al., 2006). In this group adult proliferation has been detected in the telencephalon, preoptic region, thalamus, hypothalamus, midbrain and cerebellum (Chetverukhin & Polenov, 1993; Dawley, Fingerlin, Hwang, John, & Stankiewicz, 2000; Polenov & Chetverukhin, 1993; Raucci et al., 2006). In teleost fishes, neurogenesis is abundant in all the brain extension from rostral to caudal with more than 10 neurogenic regions (the olfactory bulb; telencephalon; thalamus; epithalamus; preoptic region; hypothalamus; tectum; cerebellum; rhombencephalon; and spinal cord) identified and approximately 100 neurogenic

areas in *Apteronotus leptorhynchus* (Ekström, Johnsson, & Ohlin, 2001; Grandel, Kaslin, Ganz, Wenzel, & Brand, 2006; Meyer, 1978; Zikopoulos, Kentouri, & Dermon, 2000; Zupanc, Hinsch, & Gage, 2005; Zupanc & Horschke, 1995).

As we can become aware in these brief summary across taxonomy there are large differences in neurogenic potential between “lower” and “higher” vertebrates. In “higher” vertebrates, neurogenic regions are restricted to telencephalon and, as we came across taxonomy, more brain areas with neurogenic potential emerge and the “neurogenic brain” is achieved in fishes, the lower vertebrate of the chordata phyla.

The adult proliferation pattern in the teleost brain has been studied in detail only in three species: the gymnotiform weakly electric fish, *Apteronotus leptorhynchus*; the gasterosteiform three-spined stickleback, *Gasterosteus aculeatus*; and the cypriniform zebrafish, *Danio rerio* (Ekström et al., 2001; Grandel et al., 2006; Zupanc et al., 2005; Zupanc & Horschke, 1995). Despite the phylogenetic distance between these groups, the distribution of the new generated cells within the proliferative zones seems to be well conserved across species. In the present study we present a detailed characterization of the neurogenic areas in another teleost fish, the Perciform, Mozambique tilapia, *Oreochromis mossambicus* (Peters, 1852; teleostei, Cichlidae) including a detailed mapping of the proliferations zones in the entire brain. The results of this investigation will bring new insights to a comparative approach on fish neurogenesis since will allow us to compare the mitotic areas in a perciform, with fish from distant superorders and see how preserved these proliferative zones are.

## **Materials and methods**

### **Animals**

A total of three tilapia, *O. mossambicus* were used in this study. The fish were kept in 300 L tanks under a 12-h light/12-h dark photoperiod at a temperature of approximately 26-28°C, pH values around 7.3, and a water conductivity of 500 µS/cm. Two of these fish were female and

one was a male. In females, the total length was 10.5 cm and 11.6 cm respectively, and 17, 0 g and 18.3 g for body weight. The male had of total length of 14.6 cm and 36.8 g of body weight. The relative gonadal weight (= fresh weight of gonads divided by body weight) was 0.0476 and 0.0513 for females and 0.0115 for the male. All of the individuals were mature fish, as determined by post-mortem gonadal inspection and as indicated by the relative gonadal weight.

### **Intraperitoneal injection of 5-bromo-2'-deoxyuridine**

For labeling of S-phase cells, the fish were anesthetized in 2% urethane (Sigma) in aquarium water and injected intraperitoneally with 50  $\mu$ L of labeling reagent [aqueous solution of 5-bromo-2'-deoxyuridine (BrdU; 3 mg/mL) and 5-fluoro-2'-deoxyuridine (0.3 mg/ mL)], as supplied with the cell proliferation kit (Amersham). After a recovery period in oxygenated aquarium water, the fish were transferred to isolation tanks.

### **Intracardial perfusion and tissue preparation**

After 2 hours post administration of BrdU, the fish were deeply anesthetized by immersion into a 1.5% solution of ethyl 3-aminobenzoate methanesulfonate salt ('MS-222'; Sigma) and intracardially perfused with a flush solution containing 0.2 M phosphate buffer (PB; pH 7.4), 0.9% NaCl, 5 mg/100 mL heparin sodium salt (research grade, 150,000 IU/g; AppliChem), and 1 mL/100 mL 2% lidocaine. After all blood had been washed out, the perfusion was continued for 30 min with 2% paraformaldehyde (AppliChem) in 0.2 M PB. The brains were removed from the skull, postfixed in fixative solution for 1h at 4°C, and cryoprotected in 1 M sucrose in phosphate-buffered saline (PBS) overnight at the same temperature. After embedding the brains in a 1:1 mixture of AquaMount (Lerner Laboratories) and Tissue Tek O.C.T. Compound (Sakura), 16- $\mu$ m thick coronal sections were cut on a cryostat and mounted onto gelatin/chrome-alum-coated slides.

## **BrdU immunohistochemistry**

For detection of BrdU-labeled cells, the sections were dried, washed with 0.1 M Tris-buffered saline (TBS; pH 7.5) and incubated in 50% formamide at 65°C for 2 h. After one rinse with 2x SSC for 30 min, the sections were incubated in 2 M HCl at 37°C for 40 min. The acid was neutralized by two rinses for 10 min each in 0.1 M borate buffer, pH 8.5, followed by six rinses in TBS for 20 min each. After blocking the slides for 1 h in TBS containing 1% bovine serum albumin (BSA; fraction 5, pH 5.2; AppliChem), 1% teleostean gelatin (Sigma), 3% normal sheep serum (Sigma), and 0.3% Triton X-100 (AppliChem) (here referred to as 'Blocking Solution 1') at room temperature (RT), the sections were incubated overnight at 4°C with rat-anti-BrdU primary antibody (Oxford Biotechnology Cat. No. OBT0030), diluted 1:200 in Blocking Solution 1. The sections were washed three times for 10 min with TBS, blocked for 30 min with Blocking Solution 2 (which was identical to Blocking Solution 1, except that instead of normal sheep serum, normal donkey serum was used) and incubated for 90 min at RT with secondary antibody, Cy3-conjugated donkey anti-rat IgG (H+L) (Dianova; Cat. No. 712-165-150), diluted 1:1000 in Blocking Solution 2. Following three washes for 10 min each in TBS and three washes of 3 min in 0.1 M PBS, pH 7.4, the sections were counterstained with DAPI (2µg/ml) for 3 min at RT and then rinsed in PBS three times for three minutes each. Slides were embedded in polyvinyl alcohol containing *n*-propyl gallate.

## **Microscopic examination and data analysis**

The sections were examined under a Zeiss Axioskop epifluorescence microscope (Carl Zeiss). Brain areas were identified using the DAPI counterstaining and a partial tilapia brain map (Pepels, Meek, Bonga, & Balm, 2002). The brain sections were drawn by means of a camera lucida, and the images were scanned using a HP scanjet 7400C and further processed in Corel Draw 11. For confocal microscopy, a Zeiss LSM 510 META laser scanning microscope was used. Optical sections were taken using 10x and 20x objectives, with a pinhole opening of one Airy at a resolution of 1.024 x 1.024 pixels, using LSM5 (version 3.2; Carl Zeiss) software.

## **Results**


### **Proliferation zones**

Intraperitoneal injection of the thymidine analogue BrdU, followed by a survival time of 2 hours, revealed BrdU-labeled cells in numerous brain regions throughout the neuraxis (Figs. 1, 2; Table 1). In the following, we will describe those among these brain regions that displayed relatively high areal densities of labeled cells, compared to neighboring regions, and thus qualify to be called 'proliferation zones.'


### **Olfactory bulb**

In the olfactory bulb, BrdU labeling exhibited a differential distribution: the vast majority of the labeled cells were located in the glomerular layer ( $BO_{gl}$ ), whereas rather few cells were observed in the granular layer ( $BO_{gra}$ ) (Figs. 1a, 3a). Within the glomerular layer, labeled cells tended to occupy ventral and lateral regions. Typically, the labeled nuclei were ovoid, with the major axis (10-13  $\mu\text{m}$ ) displaying roughly twice the length of the minor axis (3-8  $\mu\text{m}$ ).


a




b




c




d



e



f



1000 μm

Fig 1 - Distribution and semi-quantitative analysis of proliferation zones in the forebrain (a-c) and midbrain (d-f) of Mozambique tilapia. Mitotic zones were revealed by the analysis of BrdU labeling in coronal sections at a post-administration survival time of 2 hours. Proliferative areas are indicated by black dots. Each dot represents approximately 50 BrdU-labeled cells.




Fig 2 - Distribution and semi-quantitative analysis of proliferation zones in the hindbrain of Mozambique tilapia. Mitotic zones were revealed by the analysis of BrdU labeling in coronal sections at a post-administration survival time of 2 hours. Proliferative areas are indicated by black dots. Each dot

represents approximately 50 BrdU-labeled cells.

Table 1- Quantitative analysis of the proliferation zones in all brain regions of tilapia. Proliferative zones were revealed by the analysis of BrdU labeling at 2 hours survival time.

<b>Number of new generated cells after 2h survival time</b>		
<b>Brain region</b>	<b>Mean</b>	<b>Range</b>
<b>Telencephalon</b>		
DMa	401	(296 - 589)
DA	184	(96 - 228)
Dla	295	(224 - 385)
tolfm	28	(32 - 52)
BO <sub>gra</sub>	20	(9 - 34)
Bogl	118	(95 - 148)
DMdd	185	(167 - 204)
DLd	148	(95 - 204)
DLp	223	(160 - 257)
Dp	87	(64 - 100)
DLv	27	(23 - 30)
DD	72	(41 - 107)
DMvv	11	(3 - 14)
DMv	60	(19 - 112)
DMvd	29	(16 - 39)
DMdv	10	(2 - 29)
VVm	96	(36 - 161)
Vd	153	(75 - 226)
DCm	2	(2 - 4)
DCad	3	(3- 6)
Vs	128	(51 - 205)
DCm	2	(2 - 4)
Dcad	3	(3- 6)
<b>Diencephalon</b>		
PPm	36	(52 - 55)
PPa	110	(67 - 162)
PPp	11	(12 - 20)
TL	27	(9 - 62)
TLAi	7	(4 - 12)
TGNI	13	(5 - 24)
ILdl	38	(21- 68)
ILvm	14	(12- 30)

PGm	10	(5 - 18)
LH	24	(1 - 61)
HV	12	(2 - 34)
TA	53	(2 - 120)
VM	18	(6 - 32)
<hr/>		
A	56	(25 - 94)
nRP	65	(46 - 82)
nRLm	237	(145 - 354)
TPP	237	(195 - 264)
CP	22	(20 - 25)
DP	20	(6 - 40)
G	47	(30 - 63)
DFld	20	(16 - 26)
DFlv	15	(7 - 27)
nRLl	265	(249 - 283)
DFm	36	(15 - 51)
TP	19	(3 - 51)
CM	13	(7 - 20)
TP	19	(3 - 51)
nX	25	(16 - 37)
PGm	10	(5 - 18)
Ch	50	(31 - 60)
H	58	(40 - 92)
DFI	30	(13 - 47)
Sc	49	(29 - 119)
CE	13	(7 - 17)
<b>Mesencephalon</b>		
<hr/>		
TeO layer 1	2176	(1845 - 2665)
TeO layer 2	82	(72 - 101)
TeO layer 3	72	(62 - 77)
TeO layer 4	106	(82 - 121)
TeO layer 5	117	(87 - 140)
TeO layer 6	186	(169 - 215)
TeO layer 7	54	(37 - 74)
tOd	41	(19 - 66)
tOv	39	(28 - 59)
TS	247	(222 - 295)
TL	300	(163 - 455)
tO	102	(80 - 131)
<b>Rombencephalon</b>		

Val <sub>gra</sub>	30	(11 - 45)
GC	342	(217- 503)
Vam <sub>mol</sub>	1285	(847 - 1769)
nIII	5	(1- 8)
MLF	4	(5 - 2)
Ipn	5	(4 - 11)
Val <sub>mol</sub>	1797	(1236 - 2478)
CCe <sub>mol</sub>	5090	(4474 - 6165)
CCe <sub>gra</sub>	575	(181 - 1171)
mol	827	(725 - 1023)
LCe <sub>gra</sub>	775	(550 - 1090)
CC	120	(116 - 125)
EG	39	(36 - 46)
LVII	52	(42 - 68)
nVd	32	(10 - 86)
MLF	52	(31- 65)
LX	238	(122 - 411)
CVEn	12	(9 - 15)
Z	141	(77 - 184)
M	569	(440 - 800)
S	100	(38 - 158)

## Telencephalon

The rostral beginning of the dorsal telencephalon was distinguished by a high areal density of BrdU-labeled cells in a narrow zone at the ventricular wall of the anterior subdivision of the dorsomedial telencephalon (DMa) (Figs. 1a, 3b). These cells stretched throughout the rostro-caudal extent of this structure. Further caudally, the density of the cells in DMa decreased, while at the dorsal aspects of this structure BrdU-labeled cells became contiguous with labeled cells that appeared in the area just beneath the pial surface of the dorsal part of the dorsal subdivision of the dorsomedial telencephalon (DMdd), the anterior part of the dorsal telencephalon (DA), the dorsal subdivision of the dorsolateral telencephalon (DLd), the ventral subdivision of dorsolateral telencephalon (DLv), and the anterior subdivision of the dorsolateral telencephalon

(DLa) (Fig. 1b). The density of labeled cells in DLv and in the dorsal part of DLa was low, relative to the densities observed in DMdd, DA, DLd, and the ventral part of DLa.

The morphology of the labeled nuclei in the different regions of the dorsal telencephalon was rather uniform, being composed of two populations. One consisted of predominantly round nuclei, with diameters of approximately 6-7  $\mu\text{m}$ , while the nuclei of the other population displayed an ovoid profile, with the lengths of the major axis and minor axis ranging between 7 and 10  $\mu\text{m}$ , and 5 and 7  $\mu\text{m}$ , respectively.

In the ventral telencephalon, the most rostrally located structure that displayed BrdU-labeled cells was the medial part of the ventral subdivision of the ventral telencephalon (Vvm) (Figs. 1b, 3c). At its rostral pole, the areal density of BrdU-labeled cells was remarkably high, but gradually decreased at more caudal levels. The labeled nuclei in Vvm were predominantly round, with diameters of 7-8  $\mu\text{m}$ .

Two other structures within the ventral telencephalon with relatively large numbers of BrdU-labeled cells were the dorsal part of the ventral telencephalon (Vd) (Figs. 1b, 3d) and the supracommissural part of the ventral telencephalon (Vs) (Fig. 1c). The labeled cells in Vs appeared as a continuation of the labeling in Vd at the level of the anterior commissure. The populations of labeled nuclei differed somewhat between the two structures. Vd exhibited mostly ovoid nuclei (major axis approximately 10  $\mu\text{m}$ , minor axis approximately 5  $\mu\text{m}$ ), whereas Vs contained, in addition to the ovoid nuclei, also round nuclei with diameters typically ranging between 7 and 8  $\mu\text{m}$ .




Fig 3 - Confocal images of BrdU-labeled cells in coronal sections of the olfactory bulb and telencephalon in adult tilapia at 2 hours post-administration survival time. a: Olfactory bulb (BO). Relatively few labeled cells were found in this brain region. Most of these cells were located in the glomerular layer ( $BO_{gl}$ ), whereas only a few were observed in the granular layer ( $BO_{gra}$ ). The labeled cells in  $BO_{gl}$  tended to occupy ventral and lateral regions (arrowhead). b: Anterior subdivision of the dorsal telencephalon (DMa). This brain region was distinguished by a high density of BrdU-labeled cells in a narrow zone at the ventricle (V). c: Medial part of the ventral subdivision of the ventral telencephalon (VVm). The majority of labeled cells were concentrated at the rostral pole of this structure, from which the section was taken. DLa, anterior subdivision of the dorsolateral telencephalon. d: Dorsal part of the ventral telencephalon (Vd). The labeled cells in this brain area are exclusively found in the immediate vicinity of the ventricle. DMvv, ventral subdivision of the ventral part of the dorsomedial telencephalon.

## Diencephalon

At the transition of telencephalon and diencephalon, BrdU labeling was most conspicuous in the anterior part of the periventricular preoptic nucleus (PPa), where labeled cells could be found over the entire dorsoventral extent (Figs. 1c, 4a). Typically, their nuclei were round, with diameters of 7-10  $\mu\text{m}$ .

### Periventricular nucleus of the posterior tuberculum

In the periventricular nucleus of the posterior tuberculum (TPP), labeled cells were restricted in medio-lateral direction to a narrow zone, less than 50  $\mu\text{m}$  wide, in the immediate vicinity of the wall of the third ventricle, but occurred throughout the dorso-lateral extent, spanning approximately 1000  $\mu\text{m}$ , of this brain nucleus (Figs. 1e, 4b). Within this area, labeled cells displayed a remarkably high packing density. Most of the labeled nuclei were round or ovoid and had typically diameters of 7-10  $\mu\text{m}$  and lengths of the major and minor axes of 7-10  $\mu\text{m}$  and 5-7  $\mu\text{m}$ , respectively.




Fig 4 - Confocal images of BrdU-labeled cells in coronal sections of the diencephalon in adult tilapia at 2

hours post-administration survival time. a: Anterior part of the periventricular preoptic nucleus (PPa). BrdU labeled cells were found over the entire dorsoventral extent of the nucleus. b: Periventricular nucleus of the posterior tuberculum (TPP). Throughout this area, labeled cells displayed a remarkably high packing density. CP, central posterior thalamic nucleus.

## Optic tectum

The optic tectum (TeO) was characterized by a large number of BrdU-labeled cells (Figs. 1d-f; 2a, b). The majority of these cells were concentrated in two areas. The first concentration of cells was observed at the dorsomedial end of the innermost layer (Layer 1) (Figs. 1e, 5a). There, Layer 1 intrudes the dorsolateral aspect of the torus longitudinalis (TL) in a conspicuous way. The second concentration of labeled cells, distinguished by a high areal density, was found in Layer 1 at the caudal pole of the tectum (Figs. 2b, 5b). In the remaining six layers of TeO (Layers 2-7), the areal density of cells was rather low, and no marked differential distribution of cells between these layers, or within the individual layers, could be detected. In the proliferation zone at the caudal pole of the tectum, a variety of nuclear morphologies were encountered, including round, ovoid, and elongated nuclei. Typically, the round nuclei had diameters of 7-10  $\mu\text{m}$ . Many of the elongated cells were remarkably slim, with the major axis (approximately 10  $\mu\text{m}$ ) often showing four times the length of the minor axis (approximately 2-3  $\mu\text{m}$ ). By contrast, in Layers 2-7 most of the cells were round (diameter approximately 7-10  $\mu\text{m}$ ).




Fig 5 - Confocal images of BrdU-labeled cells in coronal sections of the optic tectum (TeO) in adult tilapia at 2 hours post-administration survival time. a: Rostral level of TeO. Most of the new cells were concentrated in the innermost layer (Layer 1). This layer intrudes the dorsolateral aspect of the torus longitudinalis (TL). The site of the intrusion is distinguished by the presence of a dense cluster of labeled cells (arrowhead). b: Caudal level of TeO. The vast majority of the new cells were concentrated, at a uniform and high areal density, in Layer 1.

### Nucleus recessi lateralis

In the lateral part of the nucleus recessi lateralis (nRLl), a high areal density of BrdU-labeled cells was found (Figs. 1e, f). Most of these cells were located at the ventral and ventrolateral aspects of this structure (Fig. 6a). Typically, labeled nuclei were predominantly ovoid, with the major axis of approximately 10  $\mu\text{m}$  and the minor axis of 4-7  $\mu\text{m}$ .

Labeled cells also occurred at relatively high areal densities in the medial part of the nucleus recessi lateralis (nRLm), where they were concentrated in the immediate vicinity of the ventricular wall (Figs. 1e, 6b). In contrast to nRLl, in nRLm two major morphologies of labeled cells were encountered - one displaying a round shape of the nucleus (diameter 7-8  $\mu\text{m}$ ), whereas the other was characterized by an ovoid nuclear profile (major axis approximately 10  $\mu\text{m}$ ; minor axis approximately 5  $\mu\text{m}$ ).




Fig 6 - Confocal images of BrdU-labeled cells in coronal sections of the recessus lateralis (RL) in adult tilapia at 2 hours post-administration survival time. a: Lateral part of nucleus recessi lateralis (nRLl). A high areal density of BrdU-labeled cells was found at the ventral and ventrolateral aspects of this structure. b: Medial part of the nucleus recessi lateralis (nRLm). Labeled cells were equally distributed throughout the medial border of this structure. ILvm, ventromedial subdivision of the inferior lobe of the hypothalamus.

## **Cerebellum**

The largest number of BrdU-labeled cells, relative to all other brain structures, was found in the three subdivisions of the cerebellum - the valvula cerebelli, the corpus cerebelli, and the lobus caudalis.

### **Valvula cerebelli**

The valvula cerebelli consists of two parts, the lateral part (Val) and the medial part (Vam). Each of these two parts can be subdivided into a molecular layer (Val<sub>mol</sub> and Vam<sub>mol</sub>, respectively) and a granular layer (Val<sub>gra</sub> and Vam<sub>gra</sub>, respectively).

#### **Lateral part of the valvula cerebelli**

In the lateral part of the valvula, a huge number of BrdU-labeled cells were present, and almost all of them were found in the molecular layer (Figs. 2a, 7a). These cells were largely restricted to the dorsal and medial aspects of the molecular layer (Fig. 7a'), with one exception. From the dorsal area, a stream of labeled elongated cells emerged and traveled in ventral direction through the molecular layer to the associated granular layer (Fig. 7a''). Virtually all labeled nuclei were aligned parallel to the main direction of this stream. These cells were elongated, with the major axis (10-20  $\mu\text{m}$ ) of the nuclei typically being 2-3 times longer than the minor axis (approximately 5  $\mu\text{m}$ ). The remaining cells in the molecular layer of the lateral part of the valvula formed a mixed population, consisting of both round cells (diameter of the nucleus approximately 5-10

$\mu\text{m}$ ) and ovoid cells (major axis of the nucleus: approximately 10  $\mu\text{m}$ ; minor axis of the nucleus: approximately 5  $\mu\text{m}$ ).

### **Medial part of the valvula cerebelli**

Like in the lateral part of the valvula, in the medial part of the valvula virtually all BrdU-labeled cells were located in the molecular layer, and only a very few cells were observed in the granular layer (Figs. 2a, 7b). Most of the labeled cells within the molecular layer were concentrated in a region stretching up to 150  $\mu\text{m}$  laterally from the midline (Fig. 7b'). A distinct feature of the BrdU-labeled nuclei was their elongated shape. Typically, the major axis (10-20  $\mu\text{m}$ ) was 2-3 times longer than the minor axis (3-8  $\mu\text{m}$ ). Overall, the elongated nuclei displayed a non-random orientation, with the major axis of many nuclei being perpendicular to the midline.

### **Corpus cerebelli**

The corpus cerebelli (CCe) extends from the caudal pole of TeO to the caudal end of the rhombencephalon and thus covers roughly half of the dorsal surface of the brain. Like the valvula, CCe is distinguished by its molecular layer (CCe<sub>mol</sub>) and granular layer (CCe<sub>gra</sub>).

At the rostralmost levels of CCe, the vast majority of labeled cells were found in the dorsal molecular layer (Figs. 2b, c; 7c). These cells were spread over relatively wide areas within this region, exhibiting somewhat higher densities at the midline and immediately below the pial surface. More caudally, these cells became restricted to two major areas. The first area was situated below the dorsal pial surface, stretching in each hemisphere approximately 500  $\mu\text{m}$  laterally from the midline. The second area was defined by a stream of cells at the midline in the dorsal molecular layer, running from the pial surface in ventral direction until the dorsal tip of the granular layer was reached (Fig. 7c'). These labeled cells formed two populations, defined by their ovoid and elongated nuclear profiles, respectively. The ovoid cells displayed lengths of the major and minor axes of approximately 10  $\mu\text{m}$  and 5  $\mu\text{m}$ , respectively. The nuclei of the elongated cells were characterized by a length of the major axis of approximately 15  $\mu\text{m}$  and of the minor axis of approximately 5  $\mu\text{m}$ . The cells of each of the two populations exhibited a non-

random orientation, with the major axis of the nucleus predominantly parallel to the pial surface in case of the ovoid cellular population, and parallel to the midline in case of the elongated cellular population. The relatively few labeled cells in the granule layer were observed throughout this structure, but they were more concentrated in the region of the midline. Their nuclear profiles were rather uniform, with the major axis of approximately 10  $\mu\text{m}$  and the minor axis of approximately 5  $\mu\text{m}$ .

### **Lobus caudalis**

The lobus caudalis (LCe) consists of a granular layer dorsal to the rhombencephalic ventricle. In the rostral half of LCe, where the granular layer is restricted to a small area at the dorsolateral corner of the rhombencephalic ventricle, BrdU-labeled cells were restricted to this area, and their number was very low (Fig. 2c). In the caudal half, where the granular layer of LCe enlarges by an expansion in both lateral and dorsal direction, the labeled cells were distributed over the entire area defined by the granular layer, and their numbers increased markedly, compared to the numbers encountered at more rostral levels. Most of the labeled cells were either round, with typical diameters of 7-8  $\mu\text{m}$ , or ovoid, with typical lengths of the major axis of 7-12  $\mu\text{m}$  and of the minor axis of 5-7  $\mu\text{m}$ .

Dorsally to LCe, a molecular layer is situated which is bordered dorsally by CCE<sub>gra</sub> and laterally by the eminentia granularis (EG). Its association with a specific granular layer is unclear and we, therefore, refer to this as 'mol'. Within this molecular layer, a high areal density of labeled cells were found at the midline, stretching throughout its dorsoventral extent (Fig. 2c). Most of the cells were ovoid (major axis, 10  $\mu\text{m}$ ; minor axis, 7-8  $\mu\text{m}$ ).




Fig 7- Confocal images of BrdU-labeled cells in coronal sections of the cerebellum in adult tilapia at 2 hours post-administration survival time. a: Lateral part of the valvula cerebelli (Val). The majority of the new generated cells were located in the molecular layer of Val (Val<sub>mol</sub>). By contrast, in the granular layer

of this structure ( $Val_{gra}$ ) very few cells were observed. The distribution of the cells in  $Val_{mol}$  was restricted to the dorsal and medial aspect of this structure, with the cells being mostly round and ovoid. Some of these cells, indicated by arrow, are shown at higher magnification in a'. At the lateral aspect of  $Val_{mol}$ , a stripe of elongated cells was found along the dorsoventral extent. Some of these cells, indicated by arrowhead, are shown at higher magnification in a". b: Medial part of the valvula cerebelli ( $Vam$ ). Virtually all of the BrdU-labeled cells were located in the molecular layer of  $Vam$  ( $Vam_{mol}$ ). Their nuclei displayed an elongated morphology and were orientated parallel to the midline. Part of these cells are shown at higher magnification in b'. c: Corpus cerebelli ( $CCe$ ). The majority of labeled cells were found in the molecular layer of this structure ( $CCe_{mol}$ ) immediately beneath the pial surface. They stretched over several hundreds of micrometers laterally in each hemisphere. In addition, BrdU-labeled cells occurred in the midline region of the dorsal molecular layer and extended from the dorsal pial surface to the dorsal aspect of the granular layer of the  $CCe$  ( $CCe_{gra}$ ). Part of the latter cellular population, indicated by arrow, is shown in c'.

## Discussion

The present study clearly demonstrated the existence of multiples proliferation zones in the brain of the adult Mozambique tilapia. Mitotic activity was particularly pronounced in the olfactory bulb, in the Dorsal, Medial and Ventral subdivisions of Telencephalon, preoptic area and periventricular nucleus of the posterior tuberculum of the Diencephalon, optic tectum of Mesencephalon, and in the three subdivisions of the cerebellum.

Analysis of the distribution of the new generated cells shows that mitotic areas appeared normally in the vicinity of the ventricular zones. However, in actinopterygian groups, for instances teleost fish, during embryonic development telencephalon undergoes a process of eversion instead of evagination, as occur in amniote groups. This developmental difference produces a notable morphological divergence in telencephalic hemispheres, with internal ventricles in non-actinopterygian, and massive telencephalic hemispheres flanking a single ventricular cavity in actinopterygians (Northcutt, 1981). These morphological differences, together with the capability of increasing the brain weight and size throughout their lifetime, give fish the possibility to produce an incredible amount of new cells in the adult brain.

## **Cell proliferation**

Stem cells are defined by their ability to self-renew and the capacity of their progenitor to adopt multiple lineages. In addition to self-renewal, expansion of progenitor population by repeated entry into cell cycle is a fundamental feature of neurogenesis. As a result, the analysis of cell cycle activity is a critical component of neurogenesis. DNA replication during the S phase of the cell cycle has been used as an opportunity to label proliferating cells by supplying a thymidine analogue that can be incorporated into the cell (Gage, Kempermann, & Song, 2008). The thymidine analogue is retained in the post-mitotic cell to be subsequently detected in histological sections. In our study we used the 5-bromo-2'-deoxyuridine (BrdU). This thymidine analogue revealed proliferation zones in all the major subdivisions of the adult brain of tilapia, with approximately 100 proliferative areas. After 2 hours survival time, the estimated time that BrdU is metabolically available to label mitotic active cells (Zupanc & Horschke, 1995), we found a total of approximately 80,000 new cells and 91 proliferative zones (Table 1). The analysis of BrdU-positive cells also revealed different distribution patterns throughout the brain structures.

In the following section, we compare our results with findings obtained in other species. For this detailed comparison, we will focus on specific brain regions that are of special interest from a comparative point of view and were also the largest fractions of the brain with mitotic activity: olfactory bulb, dorsal telencephalon, optic tectum and cerebellum.

## **Telencephalon**

### **Olfactory bulb**

The results of the present investigation have shown that in the olfactory bulb of tilapia the majority of BrdU-labeled cells were found in the glomerular layer (BOgl). Similar results have been observed in zebrafish and gilthead sea bream (Byrd & Brunjes, 2001; Zikopoulos et al., 2000; Zupanc et al., 2005). However, in other teleost species, for instance, goldfish, mediterranean barbell, common carp and rainbow trout (Alonso, Lara, Vecino, Covenas, & Aljon, 1989), the proliferation zone was found in the innermost layer of the olfactory bulb, the

granular cell layer. These results showed that both layers, glomerular and granular respectively, have mitotic ability, and depending on the species cell proliferation can occur in the glomerular or granular cells layers.

In our study, BrdU-positive cells revealed a parallel orientation with the midline of the ventricle, what apparently indicate a migratory behaviour to the glomerular layer of the olfactory bulb. A previous study in zebrafish corroborate this idea, showing that after 4 weeks of post-BrdU administration the great majority of cells are located in the internal cellular layer (Byrd & Brunjes, 2001). In reptiles, the cells generated in the vicinity of the ventricle migrate as well into the granule cell layer (López-García et al., 1988), and in mammals part of newly generated neurons in the olfactory bulb are granule neurons, with only a small proportion acquiring a periglomerular neuronal fate (Saghatelyan, de Chevigny, Schachner, & Lledo, 2004). All this evidence suggest that that the new cells will migrate to the granular cells layer where they differentiate into granule neurons. However we need further investigation with longer survival times to confirm this hypothesis.

Another interesting thing to mention is the characteristics of the new cells. In species where the proliferation spot is the glomerular layer, the cells need to migrate to the granular cell layer and they need an appropriate stimuli and environment to induce migration. In contrast, in species where the mitotic zone is the granular cell layer, there is no need for migration or stimulation. We can hypothesize also that the brain environment and the number of generated cells is different according to the proliferation zones. We can expect higher concentration of cells in species where migration has to occur, compared to species where migration is not required, given that during the migration process, a high number of cells normally died.

Nevertheless, as the olfactory bulb is one of the two brain regions where adult neurogenesis has been described in mammals, it is especially interesting to study this phenomenon in more detail.

## **Dorsal and ventral telencephalon**

The results of 2 hours post-BrdU administration experiments in telencephalon areas were remarkably similar to those found in other teleost fishes (Ekström et al., 2001; Zikopoulos et al., 2000; Zupanc et al., 2005; Zupanc & Horschke, 1995). For tilapia, we observed a high proliferative activity in the ventricular zone. This proliferation initiated at the rostral beginning of the dorsal telencephalon (DA), and became contiguous with the dorsomedial (DMa, DMdd), and dorsolateral telencephalon (DLa, DLd, DLv and DLp), covering all the dorsal telencephalic areas.

In ventral telencephalon (Vvm, Vd and Vs), the density of BrdU-labeled cells was remarkably high. These mitotic areas, in dorsal and ventral telencephalon, correspond very well to those observed for the brown ghost, stickleback, zebrafish, salmon and gilthead sea bream (Ekström et al., 2001; Lema, Hodges, Marchetti, & Nevitt, 2005; Zikopoulos et al., 2000; Zupanc et al., 2005; Zupanc & Horschke, 1995).

The uniformity of the results obtained across different teleost species shown that mitotic activity is extremely conserved across fish taxonomy, and if we go further to other vertebrate groups, i.e. mammals, birds, reptiles and amphibians, we can also observe that telencephalic subdivisions are the only brain regions where constitutive adult neurogenesis has been well-documented, indicative of the significance of mitosis in this brain region (Kaslin, Ganz, & Brand, 2009).

Several studies showed the importance of adult neurogenesis in the telencephalon. In mammals, the existent data suggest that neurogenesis and neuronal replacement are related with the acquisition of new information (Ming & Song, 2005). In birds, this phenomenon seems to be correlated as well with memory formation. Several studies of food-hoarding, social change, and reproductive cycle, indicate that the number of new neurons increases markedly as the memory load increases (Barnea, 2009). On the basis of neuroanatomic evidence and results of functional studies (Northcutt, 2008; Wullimann & Mueller, 2004), the dorsolateral telencephalon in fish is thought to be homologous to the mammalian hippocampus, and the dorsomedial telencephalon

homologous to the amygdala, both areas associated with memory formation (Broglia et al., 2005).

Despite the fact that in fish there is no study that correlates directly cell proliferation and memory formation, there are several works that indicate such correlation. Initial studies in rainbow trout (*O.mykiss*) showed that the telencephalon area is significantly reduced in fish reared in hatcheries compared to those reared in the wild (Marchetti & Nevitt, 2003), and Lema *et al* (2005) showed that different environments induce differences in mitotic activity rates in coho salmon (*Oncorhynchus kisutch*). These results suggest that the environment plays a key role in the regulation of brain development in adult fishes, *via* changes in mitotic activity.

A considerable amount of experimental data, also involves the dorsolateral telencephalon in allocentric spatial memory (Rodriguez et al., 2002), a process that produces a brain representation of the spatial environment with geometric relationships independent of the subject own position (O'keefe & Nadel, 1979). In goldfish, a significant increase in protein synthesis in the dorsolateral telencephalon neurons was observed in fish trained in a spatial learning task, and lesions in the same region induced a dramatic impairment in place learning (Vargas, Rodriguez, Lopez, Arias, & Salas, 2000).

Based on the facts presented above, we can infer that also in fishes the formation of new cells has a relation with the acquisition of new information, as we can state for the so-called “higher” vertebrates. However, further studies have to be conducted in this direction.

## **Optic tectum**

Our findings revealed that the higher concentration of labeled cells in the optic tectum was found in Layer 1, at the caudal pole of the tectum, and once more our results are consistent with previous results published for other teleost species, such as zebrafish (Marcus, Delaney, & Easter, 1999; Zupanc et al., 2005), Goldfish (Meyer, 1978; Raymond & Easter, 1983), brown ghost (Zupanc & Horschke, 1995), gilthead sea bream (Zikopoulos et al., 2000) and stickleback (Ekström et al., 2001).

The optic tectum, the homologous structure of the superior colliculus in mammals, is characterized by a specialized cytoarchitecture and by abundant connectivity with other sensory and motor centers (Vanegas, 1984). Its general organization and function have been conserved across phylogeny in minimum details, such as the pattern of intrinsic and extrinsic connectivity and the tectoreticular projections (Torres, Pérez-Pérez, Herrero, Ligerio, & Nunez-Abadesa, 2002).

The analysis of adult neurogenesis in the visual system covered so far a broad range of non related teleost species and a similar pattern emerge among them. Germinal cells are normally absent at the rostral aspect of the tectum, and their density increases in caudal direction on both the dorsomedial and ventrolateral edges of this structure. The two arms of the tectum joined at the caudal pole of this structure, where the density of mitotic cells reaches maximum values (Raymond & Easter, 1983). This pattern of cell birth seems to be related with developmental and organization features of the optic tectum. During embryonic development, this brain structure has a pronounced rostral-to-caudal gradient of maturation, such that the first cells to differentiate are at the rostralateral pole and the last ones are at the caudomedial edge (Cowan, Martin, & Wenger, 1968).

The data from embryonic development correspond to the matrix of proliferation in the adult brain, indicating that the caudal pole of this structure retain the ability of mitotic division, as we can confirm once more with the present results for tilapia. Kirsche (1967, *in* Raymond & Easter, 1983) also propose that the proliferation zones in adult animals exist within brain areas that are the last to develop or their development persist within the brain maturation process. Regarding to organizational aspects, the continual integration of new neurons to the visual system must preserve the topography of retinal projection (Straznicky & Gaze, 1971), therefore the caudal pole of the tectum is the finest place for this renewal. These results together suggest that apart from the ecological and ethological constraints among the different species, the commonality of the proliferation zones in visual and nonvisual (electric fish) species does not appear to be related with visual inputs because the proliferation areas are constant between species.

## Cerebellum

Among all the brain regions studied for tilapia, the main cell production was found in the three subdivision of the cerebellum: valvula cerebelli, corpus cerebelli, and the lobus caudalis. An identical result was obtained in all other teleost species examined in detail, including guppy (Kranz & Richter, 1970) brown ghost (Zupanc & Horschke, 1995), gilthead sea bream (Zikopoulos et al., 2000), three-spined stickleback (Ekström et al., 2001) and zebrafish (Zupanc et al., 2005).

Beyond the proliferation zones, the distribution of new cells seems to be conserved among the different species. In all the species studied so far, the majority of new cells for corpus and valvula cerebelli, was generated in the molecular layers. For tilapia, in the dorsal aspect of corpus cerebelli immediately below the dorsal pial surface, the new cells stretched in each hemisphere approximately 500  $\mu\text{m}$  laterally from the midline. This same result was found in zebrafish, where the cells stretched 100  $\mu\text{m}$  laterally in either hemisphere (Zupanc et al., 2005), in brown ghost cells stretching up to 200-300  $\mu\text{m}$  laterally on each side of the brain (Zupanc & Horschke, 1995) and in the stickleback (Ekström et al., 2001). The  $\mu\text{m}$  of stretching are possibly correlated with the brain size and almost no variability can be found.

In valvula cerebelli, the general pattern of proliferation was largely restricted to the dorsal and medial aspects of the molecular layer. From the dorsal area, a stream of elongated cells emerged traveling in ventral direction to the associated granular layer. The elongated morphology of these cells suggest migratory activities. This result could represent a specie-specific difference. In other teleost species, valvula cerebelli has shown consistent results in the distribution of the new cells. In brown ghost, stickleback, zebrafish and gilthead sea bream, the concentration of cells were restricted to the midline of the valvula (Zupanc et al., 2005; Zupanc & Horschke, 1995), and for the latter species, new cells could also be found in the medio-lateral axis (Zikopoulos et al., 2000). These differences in proliferative zones can be related with the differences in embryonic development of the cerebellum.

In lobus caudalis, in contrast with the others cerebellum structures, BrdU-labeled cells were distributed over the entire area defined by the granular layer. The labeled cells increased

gradually from rostral to caudal aspects of this structure and the same pattern was also described for other teleost species (Ekström et al., 2001; Zikopoulos et al., 2000; Zupanc et al., 2005; Zupanc & Horschke, 1995).

The consistence of the proliferative zones is probably related with the embryonic development. Developmental studies in trout, *Salmo gairdneri* (Pouwels, 1978), identified three matrix zones in the cerebellar wall that retain mitotic characteristics permanently, matrix M, L and P. The matrix M is located at the paramedian region of the cerebellar wall, the matrix L surrounds the lateral recessus of the fourth ventricle and the matrix P connects the left and right matrix zone L. Nevertheless, developmental studies in tilapia are necessary to confirm the matrix zones that persist from early development, and see if these matrixes are able to explain the small differences found in this particular species.

## **Conclusion**

In conclusion, we have demonstrated the presence of restricted proliferation zones in all brain subdivisions of adult Mozambique tilapia. The proliferation zones are mostly restricted to ventricular areas, as described previously for other species, and mitotic active cells were only rarely observed in deep brain tissue.

The adult proliferation zones in tilapia show once more that they are extremely well conserved. In the present study, the data obtained for tilapia (perciform) was compared with other perciforms (gilthead seabram), with cypriniformes (zebrafish and guppy), gymnotiformes (weakly electric fish), and gasterosteiformes (three-spined stickleback), and the results were consistent among all the different superorders. Even when comparisons were made with amniotic groups (reptiles, birds and mammals), these proliferative zones were still conserved. Our results suggest that the presence of mitotic activity in specific brain regions is a primitive feature that has been conserved through evolution.

## Bibliography

- Alonso, J. R., Lara, J., Vecino, E., Covenas, R., & Aljon, J. (1989). Cell proliferation in the olfactory bulb of adult freshwater teleosts. *Journal of anatomy*, 163, 15.
- Altman, J. (1969). Autoradiographic and Histological Studies of Postnatal Neurogenesis .4. Cell Proliferation and Migration in Anterior Forebrain, with Special Reference to Persisting Neurogenesis in Olfactory Bulb. *Journal of Comparative Neurology*, 137(4), 433-&.
- Alvarez-buylla, A., Ling, C. Y., & Yu, W. S. (1994). Contribution of Neurons Born during Embryonic, Juvenile, and Adult Life to the Brain of Adult Canaries - Regional Specificity and Delayed Birth of Neurons in the Song-Control Nuclei. *Journal of Comparative Neurology*, 347(2), 233-248.
- Alvarez-buylla, A., Theelen, M., & Nottebohm, F. (1990). Proliferation Hot-Spots in Adult Avian Ventricular Zone Reveal Radial Cell-Division. *Neuron*, 5(1), 101-109.
- Alvarezbuylla, A., Ling, C. Y., & Yu, W. S. (1994). Contribution of Neurons Born during Embryonic, Juvenile, and Adult Life to the Brain of Adult Canaries - Regional Specificity and Delayed Birth of Neurons in the Song-Control Nuclei. *Journal of Comparative Neurology*, 347(2), 233-248.
- Barnea, A. (2009). Interactions between environmental changes and brain plasticity in birds. *General and Comparative Endocrinology*, 163(1-2), 128-134.
- Broglio, C., Gomez, A., Duran, E., Ocana, F. M., Jimenez-Moya, F., Rodriguez, F., et al. (2005). Hallmarks of a common forebrain vertebrate plan: Specialized pallial areas for spatial, temporal and emotional memory in actinopterygian fish. *Brain Research Bulletin*, 66(4-6), 277-281.
- Byrd, C. A., & Brunjes, P. C. (2001). Neurogenesis in the olfactory bulb of adult zebrafish. *Neuroscience*, 105(4), 793-801.
- Chapouton, P., Jagasia, R., & Bally-Cuif, L. (2007). Adult neurogenesis in non-mammalian vertebrates. *Bioessays*, 29(8), 745-757.
- Chetverukhin, V. K., & Polenov, A. L. (1993). Ultrastructural Autoradiographic Analysis of Neurogenesis in the Hypothalamus of the Adult Frog, *Rana-Temporaria*, with Special Reference to Physiological Regeneration of the Preoptic Nucleus .1. Ventricular Zone Cell-Proliferation. *Cell and Tissue Research*, 271(2), 341-350.
- Cowan, W. M., Martin, A. H., & Wenger, E. (1968). Mitotic Patterns in Optic Tectum of Chick during Normal Development and after Early Removal of Optic Vesicle. *Journal of Experimental Zoology*, 169(1), 71-&.
- Dawley, E. M., Fingerlin, A., Hwang, D., John, S. S., & Stankiewicz, C. A. (2000). Seasonal cell proliferation in the chemosensory epithelium and brain of red-backed salamanders, *Plethodon cinereus*. *Brain Behavior and Evolution*, 56(1), 1-13.
- Ekström, P., Johnsson, C. M., & Ohlin, L. M. (2001). Ventricular proliferation zones in the brain of an adult teleost fish and their relation to neuromeres and migration (secondary matrix) zones. *Journal of Comparative Neurology*, 436(1), 92-110.

- Font, E., Desfilis, E., Perez-Canellas, M. M., & Garcia-Verdugo, J. M. (2001). Neurogenesis and neuronal regeneration in the adult reptilian brain. *Brain Behavior and Evolution*, 58(5), 276-295.
- Gage, F. H. (2000). Mammalian neural stem cells. *Science*, 287(5457), 1433-1438.
- Gage, F. H., Kempermann, G., & Song, H. J. (2008). Adult Neurogenesis (pp. 673). New York: Cold Spring Harbor.
- García-Verdugo, J. M., Llahi, S., Ferrer, I., & Lopezgarcia, C. (1989). Postnatal Neurogenesis in the Olfactory Bulbs of a Lizard - a Tritiated-Thymidine Autoradiographic Study. *Neuroscience Letters*, 98(3), 247-252.
- Goldman, S. A., & Nottebohm, F. (1983). Neuronal Production, Migration, and Differentiation in a Vocal Control Nucleus of the Adult Female Canary Brain. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences*, 80(8), 2390-2394.
- Grandel, H., Kaslin, J., Ganz, J., Wenzel, I., & Brand, M. (2006). Neural stem cells and neurogenesis in the adult zebrafish brain: Origin, proliferation dynamics, migration and cell fate. *Developmental Biology*, 295(1), 263-277.
- Kaslin, J., Ganz, J., & Brand, M. (2009). Proliferation, neurogenesis and regeneration in the non-mammalian vertebrate brain. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 363(1489), 101-122.
- Kranz, D., & Richter, W. (1970). [Autoradiographic studies on the synthesis of DNA in the cerebellum and medulla oblongata of teleosts of various ages]. *Z Mikrosk Anat Forsch*, 82(2), 264-292.
- Lema, S. C., Hodges, M. J., Marchetti, M. P., & Nevitt, G. A. (2005). Proliferation zones in the salmon telencephalon and evidence for environmental influence on proliferation rate. *Comparative Biochemistry and Physiology a-Molecular & Integrative Physiology*, 141(3), 327-335.
- López-García, C., Molowny, A., Garciverdugo, J. M., & Ferrer, I. (1988). Delayed Postnatal Neurogenesis in the Cerebral-Cortex of Lizards. *Developmental Brain Research*, 43(2), 167-174.
- Marchetti, M. P., & Nevitt, G. A. (2003). Effects of hatchery rearing on brain structures of rainbow trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes*, 66(1), 9-14.
- Marchioro, M., Nunes, J. M. D. M., Ramalho, A. M. R., Molowny, A., Perez-Martinez, E., Ponsoda, X., et al. (2005). Postnatal neurogenesis in the medial cortex of the tropical lizard *Tropidurus hispidus*. *Neuroscience*, 134(2), 407-413.
- Marcus, R. C., Delaney, C. L., & Easter, S. S. (1999). Neurogenesis in the visual system of embryonic and adult zebrafish (*Danio rerio*). *Visual Neuroscience*, 16(3), 417-424.
- Meyer, R. L. (1978). Evidence from Thymidine Labeling for Continuing Growth of Retina and Tectum in Juvenile Goldfish. *Experimental Neurology*, 59(1), 99-111.
- Ming, G. L., & Song, H. J. (2005). Adult neurogenesis in the mammalian central nervous system. *Annual Review of Neuroscience*, 28, 223-250.
- Northcutt, R. G. (1981). Evolution of the Telencephalon in Non-Mammals. *Annual Review of Neuroscience*, 4, 301-350.
- Northcutt, R. G. (2008). Forebrain evolution in bony fishes. *Brain Research Bulletin*, 75(2-4), 191-205.
- O'keefe, J., & Nadel, L. (1979). The Cognitive Map as a Hippocampus. *Behavioral and Brain Sciences*, 2(4), 520-528.

- Pepels, P. P. L. M., Meek, J., Bonga, S. E. W., & Balm, P. H. M. (2002). Distribution and quantification of corticotropin-releasing hormone (CRH) in the brain of the teleost fish *Oreochromis mossambicus* (tilapia). *Journal of Comparative Neurology*, 453(3), 247-268.
- Pérez-Sánchez, F., Molowny, A., Garciaverdugo, J. M., & Lopezgarcia, C. (1989). Postnatal Neurogenesis in the Nucleus Sphericus of the Lizard, *Podarcis-Hispanica*. *Neuroscience Letters*, 106(1-2), 71-75.
- Polenov, A. L., & Chetverukhin, V. K. (1993). Ultrastructural Autoradiographic Analysis of Neurogenesis in the Hypothalamus of the Adult Frog, *Rana-Temporaria*, with Special Reference to Physiological Regeneration of the Preoptic Nucleus .2. Types of Neuronal Cells Produced. *Cell and Tissue Research*, 271(2), 351-362.
- Pouwels, E. (1978). Development of Cerebellum of Trout, *Salmo-Gairdneri* .1. Patterns of Cell-Migration. *Anatomy and Embryology*, 152(3), 291-308.
- Rauci, F., Di Fiore, M. M., Pinelli, C., D'Aniello, B., Luongo, L., Polese, G., et al. (2006). Proliferative activity in the frog brain: A PCNA-immunohistochemistry analysis. *Journal of Chemical Neuroanatomy*, 32(2-4), 127-142.
- Raymond, P. A., & Easter, S. S. (1983). Post-Embryonic Growth of the Optic Tectum in Goldfish .1. Location of Germinal Cells and Numbers of Neurons Produced. *Journal of Neuroscience*, 3(5), 1077-1091.
- Rodriguez, F., Lopez, J. C., Vargas, J. P., Broglio, C., Gomez, Y., & Salas, C. (2002). Spatial memory and hippocampal pallium through vertebrate evolution: Insights from reptiles and teleost fish. *Brain Research Bulletin*, 57(3-4), 499-503.
- Saghatelian, A., de Chevigny, A., Schachner, M., & Lledo, P. M. (2004). Tenascin-R mediates activity-dependent recruitment of neuroblasts in the adult mouse forebrain. *Nature Neuroscience*, 7(4), 347-356.
- Straznicky, K., & Gaze, R. M. (1971). Growth of Retina in *Xenopus-Laevis* - an Autoradiographic Study. *Journal of Embryology and Experimental Morphology*, 26(Aug), 67-&.
- Torres, B., Pérez-Pérez, M. P., Herrero, L., Ligerroa, M., & Nunez-Abadesa, P. A. (2002). Neural substrata underlying tectal eye movement codification in goldfish. *Brain Research Bulletin*, 57(3-4), 354-348.
- Vanegas, H. (1984). *Comparative neurology of the optic tectum*. New york: Plenum Press.
- Vargas, J. P., Rodriguez, F., Lopez, J. C., Arias, J. L., & Salas, C. (2000). Spatial learning-induced increase in the argyrophilic nucleolar organizer region of dorsolateral telencephalic neurons in goldfish. *Brain Research*, 865(1), 77-84.
- Wullimann, M. F., & Mueller, T. (2004). Teleostean and mammalian forebrains contrasted: Evidence from genes to behavior (vol 475, pg 143, 2004). *Journal of Comparative Neurology*, 478(4), 427-428.
- Zikopoulos, B., Kentouri, M., & Dermon, C. R. (2000). Proliferation zones in the adult brain of a sequential hermaphrodite teleost species (*Sparus aurata*). *Brain Behavior and Evolution*, 56(6), 310-322.
- Zupanc, G. K. H., Hinsch, K., & Gage, F. H. (2005). Proliferation, migration, neuronal differentiation, and long-term survival of new cells in the adult zebrafish brain. *Journal of Comparative Neurology*, 488(3), 290-319.

Zupanc, G. K. H., & Horschke, I. (1995). Proliferation Zones in the Brain of Adult Gymnotiform Fish - a Quantitative Mapping Study. *Journal of Comparative Neurology*, 353(2), 213-233.

## **Annex I**

### **State of the art**

#### **The development of the nervous system**

The nervous system development is a complex process that includes initial expression of proneural genes and subsequent neuronal commitment, migration of these committed precursors, axonal outgrowth and guidance, and finally functional synapse formation followed by fine-tuning of connections (Demir, Singh, Klimaschewski, & Kurnaz, 2009).

Vertebrate neurogenesis is essentially regulated by proneural genes, typically basic helix-loop-helix (bHLH) proteins, such as: Neurogenin 2 (Ngn2), Math-1 (mammalian atonal homolog) or Mash-1 (mammalian achaete-Scute complex)-1 (Kim, Battiste, Nakagawa, & Johnson, 2008). All of these respond to a range of signaling molecules, such as Fibroblast Growth Factor (FGF), Bone Morphogenic Protein (BMP) / Transforming Growth Factor (TGF)- $\beta$  and Notch signaling.

The generation of distinct neuronal subpopulation from neural stem cells (NSCs) requires a specific birth order, regulated by temporal regulators, for instance Castor and Seven-up, which initially control the cell size and identity of the neuronal progeny, after they switch off the cell cycle (Maurange, Cheng, & Gould, 2008). Therefore, the regulation of cell cycle entry and exit is a critical component of neuronal development (Demir et al., 2009).

#### **The importance of the cell cycle**

Cell cycle it is a simple process that begins when a new cell is created by a mitotic division. At the beginning, the cell resides in G1 phase in a normal diploid state (2N). The cell then enters the DNA synthesis or S-phase of the cell cycle where it doubles its genome to become tetraploid (4N). After a brief pause (G2), the cell enters mitosis (M phase) where it divides to produce two daughter cells, each one in the 2N stage, the daughter cells then reenter G1. About half-way to G1, each daughter cell passes through a restriction check point at which each of the two daughter

cells decides independently whether or not it will reenter the cell cycle to repeat the process or exit the cell cycle and stop dividing (Nowakowski & Hayes, 2008). Since there are two daughter cells and two choices (reenter or exit) there are four possible combination of decision and three different types of division:

1. Two daughter cells reenter the cell cycle (*Symmetric nonterminal cell division*), producing exponential growth of the population ( $2^n$ ).
2. Neither daughter cell reenters the cell cycle (*Symmetric terminal cell division*), the two daughter cells persist postmitotic, and the proliferative lineage disappears.
3. One daughter cell reenters the cell cycle and the other exits the cell cycle (*asymmetric cell division*) producing what is called steady-state growth. The proliferating population remains at a constant size and the output is one new cell per proliferating cell (Gage, Song, & Kempermann, 2008).

The more realistic scenario it is an intermediate one, a mixture of both types of symmetric and asymmetric cell divisions. At the beginning there will be a period of symmetric cell division to produce a “founder” population. At this time, some of these cells will begin to produce daughter cells that exit the cell cycle. Thus, at least some of the cells will need to divide either asymmetrically or in a symmetric terminal way. For the next several cell cycles, the proportion of daughter cells that exit the cycle remains small, less than 50%, this means that the proportion of daughter cell that reenter the cell cycle would be greater than 50%. As long as these conditions are met, more proliferative cells will be present at the end of each cell cycle than there had been at the beginning, and thus, the proliferative population will continue to expand even as neurons are being produced. At some point, more daughter cells will exit the cell cycle than reenter, and when this point is reached, the proliferative population begins to disappear, eventually completely exhausting itself (Nowakowski & Hayes, 2008).

After exiting the cell cycle the new cells became committed with a differentiation program leading to neuronal differentiation. Exiting from the cell cycle involves an elaborated series of microtubule dynamics to generate the axonal and dendritic structures necessary for the correct functioning of the nervous system. Microtubules are highly dynamic structures within cells that allow rapid changes during processes such as cell division, migration, or differentiation

(Conde & Caceres, 2009). They are also one of the major components of the cytoskeleton. The neuronal cytoskeleton is essential for establishing the cell shape and for axonal transport.

While neurons differentiate into different subtypes they also assume a highly polar structure with a particular morphology of the cytoskeleton, thereby making it almost impossible for any differentiated cells to re-enter the cell cycle. It has been observed that dysregulated or forced cell cycle re-entry is closely linked to neurodegeneration and apoptosis in neurons (Conde & Caceres, 2009).

### **Adult neurogenesis**

The term adult neurogenesis is used nowadays to describe the formation of new neurons born from stem cells residing in discrete locations. These new cells migrate, differentiate and mature into new integrated functioning cells.

The idea that new neurons were generated in adult brains was a cause of skepticism, mainly because as we showed previously, neurons are postmitotic cells, a judgment that still holds. If neurons are postmitotic cells, how could they divide to produce new neurons? This problem was solved, with the discovery of small groups of stem cells in specific locations of the adult brain, and this finding revolutionized the neuroscience field. Since this discovery, a lot of research has been directed towards understanding the physiology and the molecular mechanisms underlying neuronal differentiation.

### **Adult neurogenesis a historical view**

Santiago Ramon y Cajal in 1913 conclude that neurons are generated exclusively during the prenatal phase of development, and since then, the “no new neurons after birth” became a central dogma in neuroscience for almost a century (Gross, 2000). The generation of new neurons in the adult brain was first reported using the  $[H^3]$ -thymidine technique in a three-day old mouse (Smart, 1961). Soon after, Altman and colleagues published a series of papers reporting evidence for new neurons in various regions of adult rats including the dentate gyrus of the hippocampus (Altman & Das, 1965) neocortex (Altman & Das, 1966) and olfactory bulb(Altman, 1969).

However, little attention was given to these studies. The issue of adult neurogenesis was revisited when in 1977 when Kaplan & Hinds (Kaplan & Hinds, 1977) demonstrated that newborn neurons in the hippocampus survived for a long period of time. These new neurons also appeared to receive synaptic inputs (Kaplan & Bell, 1983) and extend axon projections to their target area (Stanfield & Trice, 1988); they were functional cells.

The field was revolutionized by the introduction of 5-bromo-2'deoxyuridine (BrdU) a synthetic thymidine analogue as a marker of the cell cycle (Gratzner, 1982). Because BrdU can be detected by immunocytochemistry for phenotypic and quantification analysis, this approach remains the most commonly used technique in the field. Before the end of twentieth century, adult neurogenesis was observed with BrdU incorporation in all vertebrates examined, including samples from human patients (Eriksson et al., 1998).

### **Adult neurogenesis in the mammalian brain**

In mammals active neurogenesis occurs only in discrete regions of the intact adult central nervous system (CNS) (Alvarez-Buylla & Lim, 2004). From rodents to primates, neurons are generated continuously in the subventricular zone (SVZ) of the telencephalic lateral ventricle where they migrate anteriorly through the rostral migratory stream (RMS) into the olfactory bulb to become interneurons and in the subgranular zone (SGZ) of dentate gyrus of the hippocampus (Ming & Song, 2005).

These two brain regions contain a microenvironment that is permissive for neural development to occur. This microenvironment is called neurogenic “niche”. The niche comprises the precursor cells themselves, their immediate progeny and immature neurons, other glial cells and endothelia, very likely immune cells, microglia, macrophages, and an extracellular matrix. The niche is surrounded by a common basal membrane (Mercier, Kitasako, & Hatton, 2002). This combination of factors provides a unique milieu to the niche allowing neural development to occur.

The identity of neural stem cells that produce new neurons in the adult CNS has been subject of debate for the past few years, however the astrocyte hypothesis is currently the prevalent one (Alvarez-Buylla & Lim, 2004). During fetal development, astrocytes are born after most neurons and are thus unable to provide neurogenic signals for multipotent stem cells

(Temple, 2001). It becomes clear, that astrocytes in the adult CNS are not merely supporting cells as traditionally believed (Ming & Song, 2005). Just like neurons, astrocytes have a broad diversity of subtypes and functions, some behave like stem cells (Seri, Garcia-Verdugo, McEwen, & Alvarez-Buylla, 2001), some provide neurogenic signals (Lim & Alvarez-Buylla, 1999), and some provide synaptogenic factors (Song, Stevens, & Gage, 2002). The molecular mechanisms underlying neural fate specification during adult neurogenesis are just beginning to be elucidated. The fate choice is influenced by a cohort of proliferating, gliogenic, and neurogenic signals within the niche. Bone morphogenic protein (BMP) signaling, has been shown to instruct adult neural progenitors to adopt a glial fate (Lim et al., 2000). Neural differentiation from adult neural progenitors in the neurogenic niche proceeds partially because of the local presence of BMP antagonists. Ependymal cells in the SVZ secrete Noggin (Lim et al., 2000), and astrocytes in the SGZ secrete neurogenesis-1 (Ueki et al., 2003), to serve as the BMP antagonist, respectively.

There is a cohort of molecules and transcriptional factor that appears to regulate adult neurogenesis. This regulation can occur at transcriptional level or at epigenetic level both leading to neuronal differentiation. In an overall view, there are distinct steps for neural development, and these steps can be divided in four phases: a precursor cell phase, an early survival phase, a postmitotic maturation phase and a late survival phase.

In the precursor cell phase there is the expansion of the pool of cells that might differentiate into neurons. The early survival phase marks the exit from the cell cycle, most newborn cells are eliminated within days after they were born. The postmitotic maturation phase is associated with the establishment of functional connections, the growth of axon and dendrites, and synaptogenesis. The late survival phase represents a period of fine-tuning. It has been estimated that the entire period of adult neurogenesis takes approximately 7 weeks (Kempermann, Song, & Gage, 2008). After this period of time, the new neurons become indistinguishable from their older neighbors, making an integral part in the neuronal network.

From what we stated before three major principles can be applied to adult neurogenesis in mammalian brains:

1. Adult neurogenesis is a process not an event, and appears to be limited to two neurogenic regions: the hippocampus and the olfactory bulb.
2. Adult neurogenesis appears to persist throughout life, but it does not produce great number of neurons after early adulthood. It is not a mass phenomenon and appears to make a qualitative rather than a quantitative contribution.
3. The new neurons appear to make a contribution to normal brain function, in the sense of structural plasticity(Gage et al., 2008).

### **Adult neurogenesis in non-mammalian vertebrates**

A lot of research has been directed towards understanding the physiological factors and underlying molecular mechanisms that drive neural stem cell to differentiate. Despite the fact that the most prominent models in the study of adult neurogenesis are rats and mice because they are closely related to humans, neurogenesis within the mature brain is not restricted to mammals, and affects various other vertebrates to a larger extent, including birds, reptiles, amphibians and fishes. Fascinating insights have been gained from these non-mammalian species, which provide comparative models to study the function of adult neurogenesis, including both the cellular composition of the neurogenic microenvironment, and for orienting newborn neurons towards specific identities(Chapouton, Jagasia, & Bally-Cuif, 2007).

Like in the adult mammalian brain, in non-mammalian brains the majority of the mitotic cells are found in discrete areas called “proliferation zones or niches” and most of these zones are located near the surfaces of ventricles or related systems.

In birds, adult neurogenesis was found in songbird learners (canary, blackbirds, starlings, zebra finches, song sparrows, towhees (Scharff, Kirn, Grossman, Macklis, & Nottebohm, 2000), non-song-learners and in food-storing birds(Barnea & Nottebohm, 1994). The proliferative zones were found dispersed along the telencephalic lateral ventricle, particularly in the ventral and dorsal aspects of the lateral wall of this structure (Alvarez-buylla, Ling, & Yu, 1994; Alvarez-buylla, Theelen, & Nottebohm, 1990; Goldman & Nottebohm, 1983). A substantial amount of newly generated cells die, and it has been reported that two-thirds of the cells degenerate failing

the neuronal integration (Alvarez-Buylla & Nottebohm, 1988; Alvarez-buylla et al., 1990). In the hippocampus of adult birds, the recruitment of neurons makes up for net loss with no gain in the total number of neurons (Barnea & Nottebohm, 1996). Thus, the total number of neurons remains constant in the avian hippocampus. In contrast to mammals, the major destination of tangentially migrating neuroblasts, the RMS in songbirds, is the lobus parolfactorius in the basal ganglia and not the olfactory bulb (Alvarez-buylla et al., 1994). Several other telencephalic regions also receive a significant amount of newly produced neurons(Alvarez-buylla et al., 1994), like the hippocampus and striatum structures (i.e. the lobus paraolfactorius, LPO, containing Area X).

In reptiles, post-natal proliferation have been found in several brain areas such as the olfactory bulb, striatum, dorsoventricular ridge, cortex, nucleus sphericus and in a much lesser extent within the cerebellum (Font, Desfilis, Perez-Canellas, & Garcia-Verdugo, 2001; Garcia-Verdugo, Llahi, Ferrer, & Lopezgarcia, 1989; Lopez-Garcıa, Molowny, Garciaverdugo, & Ferrer, 1988; Marchioro et al., 2005; Perez-Sanchez, Molowny, Garciaverdugo, & Lopezgarcia, 1989). New cells are generated in the vicinity of the ventricle and some of the produced cells migrate along radial glia into the granule cell layer. In contrast to mammals, where differentiation takes at least 4 weeks, in reptiles only 1 to 2 weeks are needed for the cells to migrate and differentiate into neurons in the cortex of *Podarcis* and *Tropidurus* (Lopez-Garcıa, Molowny, Garcia-Verdugo, Martinez-Guijarro, & Bernabeu, 1990; Marchioro et al., 2005). This result is supported by retrograde tracings that suggest emission of proper projection pathways and formation of synapses, key signs of functional integration into the neurocircuitry (Lopez-Garcıa et al., 1990; Molowny, Nacher, & Lopez-Garcia, 1995).

In the amphibian brain, there is only one study that describes the proliferative zones (Raucci et al., 2006). In this group adult proliferation has been detected in the telencephalon, preoptic region, thalamus, hypothalamus, midbrain and cerebellum (Chetverukhin & Polenov, 1993; Dawley, Fingerlin, Hwang, John, & Stankiewicz, 2000; Raucci et al., 2006).Unfortunately, there is lack o information about adult proliferation in this group.

In teleost fishes, neurogenesis is abundant in all the brain extension from rostral to caudal with more than 10 neurogenic regions identified: (the olfactory bulb; telencephalon; thalamus; epithalamus; preoptic region; hypothalamus; tectum; cerebellum; rhombencephalon; and spinal cord) and approximately 100 neurogenic areas in *Apteronotus leptorhynchus* (Ekstrom,

Johnsson, & Ohlin, 2001; Grandel, Kaslin, Ganz, Wenzel, & Brand, 2006; Meyer, 1978; Zikopoulos, Kentouri, & Dermon, 2000; Zupanc, Hinsch, & Gage, 2005; Zupanc & Horschke, 1995). Adult neurogenesis in fish is a massive ongoing process in contrast with the qualitative process that occurs in mammals (Figure 1). In the cerebellum of *Apteronotus leptorhynchus*, quantitative analysis has shown that approximately 100 000 mitotic cells in the S-phase are found in the proliferative zone of the cerebellum of a teleost during any 2 h period (Zupanc & Horschke, 1995). After the first weeks, approximately half of the produced cells die (Soutschek & Zupanc, 1996), being the rest integrated into the network as granule neurons presumably leading to a slow increase in cerebellar size.




Figure 1 - Parasagittal schematic overviews of the adult proliferation pattern and neurogenic regions in the brain of adult vertebrates. (a) Rodent (b) Fish (zebrafish). *Adpatated from Kaslin, J. (2009).*

But why does neurogenesis cease in all except two regions of the mammalian brain shortly after birth, but persists throughout life in numerous regions of the teleost fish brain? A possible answer is given by the fact that in the majority of teleost species the body and brain size increases during most of their life time (Brandstatter & Kotrschal, 1990). Although many species of teleosts and mammals grow during postembryonic stages of development and even throughout life, there is a distinct difference in the growth pattern between these two taxonomic

classes (Zupanc, 2008). In mammals postembryonic growth is the result of an increase in size but not in number of individual muscle fibers (Rowe & Goldspink, 1969), whereas in fish, both the number of muscle fibers and the volume of individual fibers increase (Rowlerson & Veggetti 2001 in Zupanc, 2008). It is, therefore, possible that the hyperplasia of peripheral motor elements prompts a concomitant increase in the number of central neurons involved in neural control of associated muscle activity. Along similar lines, the number of sensory receptor cells, receptor organs, or receptor units in the periphery increases with age, as has been shown in several species of fish. Such a formation of new sensory elements has been demonstrated for sensory hair cells in the inner ear of sharks (Corwin, 1981), for retinal cells in the eyes of goldfish (Johns & Easter, 1977), and for electrosensory receptor organs in the gymnotiform fish *Sternopygus dariensis* (Zakon, 1984). Thus, like the formation of new motor elements, the continuous increase in the number of sensory elements in fish may lead to the generation of new neurons involved in the processing of sensory information, ensuring a numerical matching of central neurons and peripheral sensory elements (Zupanc, 2008).

The numerical matching hypothesis makes predictions about the relationship between peripheral elements, such as muscle fibers or sensory receptor cells, and central elements in the brain involved in motor control of these muscle fibers, or in processing of sensory information received through these receptors, during growth. A number of observations suggest that the ratio of the corresponding peripheral elements and central elements is highly constant and maintained when the number of peripheral elements changes. Since such growth through addition of new peripheral elements to the population of existing ones is characteristic of teleost fish, but largely absent in adult mammals, the numerical matching hypothesis can explain the difference in the extent of adult neurogenesis between teleosts and mammals.

In fish, there is some experimental evidence supporting this hypothesis. Raymond in 1983 (Raymond, Easter, Burnham, & Powers, 1983), found in goldfish that permanent removal of the optic input by enucleation of the eye results in sustained depression of mitotic activity in the tectal proliferation zone on the denervated side compared to the intact one. Temporary denervation by optic nerve crush initially has a similar effect, but upon reinnervation of the tectum by the regenerating optic fibers, proliferation is enhanced on the experimental side compared to the control side.

Comparing the three major principles applied to adult neurogenesis in mammalian brains with fish brains:

<b>Mammals</b>	<b>Fish</b>
Adult neurogenesis is a process limited to two neurogenic regions: the hippocampus and the olfactory bulb.	Adult neurogenesis is a process spread in all the brain extension from rostral to cordal
It is <u>not</u> a mass phenomenon and appears to make a qualitative contribution.	It is a mass phenomenon with a quantitative contribution.
The new neurons appear to make a contribution for structural plasticity.	The new neurons appear to make a contribution to the increasing of the brain size and also for structural plasticity in specific areas.

In the present study we present a detailed characterization of the neurogenic areas in another teleost fish, the Mozambique tilapia, *Oreochromis mossambicus* (Peters, 1852; teleostei, Cichlidae) including a detailed mapping of the proliferations zones in the entire brain. The results of this investigation will bring new insights to a comparative approach on fish neurogenesis since will allow us to compare the mitotic areas in tilapia with other fishes that were already described.

Given the enormous potential that fish have to produce new neurons and the abundance of proliferation zones in the adult brain, fishes are one of the best models to study and understand the mechanisms underlying neuronal proliferation.

*“For many problems there is an animal in which it can be most conveniently studied”*

August Krogh

## Bibliography

- Altman, J. (1969). Autoradiographic and Histological Studies of Postnatal Neurogenesis .4. Cell Proliferation and Migration in Anterior Forebrain, with Special Reference to Persisting Neurogenesis in Olfactory Bulb. *Journal of Comparative Neurology*, 137(4), 433-&.
- Altman, J., & Das, G. D. (1965). Autoradiographic and Histological Evidence of Postnatal Hippocampal Neurogenesis in Rats. *Journal of Comparative Neurology*, 124(3), 319-&.
- Altman, J., & Das, G. D. (1966). Autoradiographic and histological studies of postnatal neurogenesis. I. A longitudinal investigation of the kinetics, migration and transformation of cells incorporating tritiated thymidine in neonate rats, with special reference to postnatal neurogenesis in some brain regions. *J Comp Neurol*, 126(3), 337-389.
- Alvarez-Buylla, A., & Lim, D. A. (2004). For the long run: maintaining germinal niches in the adult brain. *Neuron*, 41(5), 683-686.
- Alvarez-buylla, A., Ling, C. Y., & Yu, W. S. (1994). Contribution of Neurons Born during Embryonic, Juvenile, and Adult Life to the Brain of Adult Canaries - Regional Specificity and Delayed Birth of Neurons in the Song-Control Nuclei. *Journal of Comparative Neurology*, 347(2), 233-248.
- Alvarez-Buylla, A., & Nottebohm, F. (1988). Migration of young neurons in adult avian brain. *Nature*, 335(6188), 353-354.
- Alvarez-buylla, A., Theelen, M., & Nottebohm, F. (1990). Proliferation Hot-Spots in Adult Avian Ventricular Zone Reveal Radial Cell-Division. *Neuron*, 5(1), 101-109.
- Barnea, A., & Nottebohm, F. (1994). Seasonal recruitment of hippocampal neurons in adult free-ranging black-capped chickadees. *Proceedings of the National Academy of Sciences of the United States of America*, 91(23), 11217-11221.
- Barnea, A., & Nottebohm, F. (1996). Recruitment and replacement of hippocampal neurons in young and adult chickadees: an addition to the theory of hippocampal learning. *Proceedings of the National Academy of Sciences of the United States of America*, 93(2), 714-718.
- Brandstatter, R., & Kotschal, K. (1990). Brain growth patterns in four European cyprinid fish species (Cyprinidae, Teleostei): roach (*Rutilus rutilus*), bream (*Abramis brama*), common carp (*Cyprinus carpio*) and sabre carp (*Pelecus cultratus*). *Brain, behavior and evolution*, 35(4), 195-211.
- Broglio, C., Gomez, A., Duran, E., Ocana, F. M., Jimenez-Moya, F., Rodriguez, F., et al. (2005). Hallmarks of a common forebrain vertebrate plan: Specialized pallial areas for spatial, temporal and emotional memory in actinopterygian fish. *Brain Research Bulletin*, 66(4-6), 277-281.
- Chapouton, P., Jagasia, R., & Bally-Cuif, L. (2007). Adult neurogenesis in non-mammalian vertebrates. *Bioessays*, 29(8), 745-757.
- Chetverukhin, V. K., & Polenov, A. L. (1993). Ultrastructural Autoradiographic Analysis of Neurogenesis in the Hypothalamus of the Adult Frog, *Rana-Temporaria*, with Special Reference to Physiological Regeneration of the Preoptic Nucleus .1. Ventricular Zone Cell-Proliferation. *Cell and Tissue Research*, 271(2), 341-350.
- Conde, C., & Caceres, A. (2009). Microtubule assembly, organization and dynamics in axons and dendrites. *Nature reviews. Neuroscience*, 10(5), 319-332.

- Dawley, E. M., Fingerlin, A., Hwang, D., John, S. S., & Stankiewicz, C. A. (2000). Seasonal cell proliferation in the chemosensory epithelium and brain of red-backed salamanders, *Plethodon cinereus*. *Brain Behavior and Evolution*, *56*(1), 1-13.
- Demir, O., Singh, S., Klimaschewski, L., & Kurnaz, I. A. (2009). From birth till death: neurogenesis, cell cycle, and neurodegeneration. *Anatomical record (Hoboken, N.J. : 2007)*, *292*(12), 1953-1961.
- Ekström, P., Johnsson, C. M., & Ohlin, L. M. (2001). Ventricular proliferation zones in the brain of an adult teleost fish and their relation to neuromeres and migration (secondary matrix) zones. *Journal of Comparative Neurology*, *436*(1), 92-110.
- Eriksson, P. S., Perfilieva, E., Bjork-Eriksson, T., Alborn, A. M., Nordborg, C., Peterson, D. A., et al. (1998). Neurogenesis in the adult human hippocampus. *Nature Medicine*, *4*(11), 1313-1317.
- Font, E., Desfilis, E., Perez-Canellas, M. M., & Garcia-Verdugo, J. M. (2001). Neurogenesis and neuronal regeneration in the adult reptilian brain. *Brain Behavior and Evolution*, *58*(5), 276-295.
- Gage, F. H., Song, H., & Kempermann, G. (2008). Adult Neurogenesis: A Prologue. In F. H. Gage, G. Kempermann & H. Song (Eds.), *Adult Neurogenesis* (pp. 3). New York: Cold Spring Harbor Laboratory Press.
- García-Verdugo, J. M., Llahi, S., Ferrer, I., & Lopezgarcia, C. (1989). Postnatal Neurogenesis in the Olfactory Bulbs of a Lizard - a Tritiated-Thymidine Autoradiographic Study. *Neuroscience Letters*, *98*(3), 247-252.
- Goldman, S. A., & Nottebohm, F. (1983). Neuronal Production, Migration, and Differentiation in a Vocal Control Nucleus of the Adult Female Canary Brain. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences*, *80*(8), 2390-2394.
- Grandel, H., Kaslin, J., Ganz, J., Wenzel, I., & Brand, M. (2006). Neural stem cells and neurogenesis in the adult zebrafish brain: Origin, proliferation dynamics, migration and cell fate. *Developmental Biology*, *295*(1), 263-277.
- Gratzner, H. G. (1982). Monoclonal antibody to 5-bromo- and 5-iododeoxyuridine: A new reagent for detection of DNA replication. *Science*, *218*(4571), 474-475.
- Gross, C. G. (2000). Neurogenesis in the adult brain: death of a dogma. *Nature reviews. Neuroscience*, *1*(1), 67-73.
- Johns, P. R., & Easter, S. S., Jr. (1977). Growth of the adult goldfish eye. II. Increase in retinal cell number. *The Journal of comparative neurology*, *176*(3), 331-341.
- Kaplan, M. S., & Bell, D. H. (1983). Neuronal proliferation in the 9-month-old rodent-radioautographic study of granule cells in the hippocampus. *Experimental brain research.*, *52*(1), 1-5.
- Kaplan, M. S., & Hinds, J. W. (1977). Neurogenesis in Adult Rat - Electron-Microscopic Analysis of Light Autoradiographs. *Science*, *197*(4308), 1092-1094.
- Kempermann, G., Song, H., & Gage, F. H. (2008). Neurogenesis in the Adult Hippocampus. In F. H. Gage, G. Kempermann & H. Song (Eds.), *Adult Neurogenesis* (pp. 161). New York: Cold Spring Harbor Laboratory Press.
- Kim, E. J., Battiste, J., Nakagawa, Y., & Johnson, J. E. (2008). *Ascl1* (*Mash1*) lineage cells contribute to discrete cell populations in CNS architecture. *Molecular and cellular neurosciences* *38*(4), 595-606.

- Lim, D. A., & Alvarez-Buylla, A. (1999). Interaction between astrocytes and adult subventricular zone precursors stimulates neurogenesis. *Proceedings of the National Academy of Sciences of the United States of America* 96(13), 7526-7531.
- Lim, D. A., Tramontin, A. D., Trevejo, J. M., Herrera, D. G., Garcia-Verdugo, J. M., & Alvarez-Buylla, A. (2000). Noggin antagonizes BMP signaling to create a niche for adult neurogenesis. *Neuron*, 28(3), 713-726.
- López-García, C., Molowny, A., Garcia-Verdugo, J. M., Martinez-Guijarro, F. J., & Bernabeu, A. (1990). Late generated neurons in the medial cortex of adult lizards send axons that reach the Timm-reactive zones. *Brain research. Developmental brain research*, 57(2), 249-254.
- López-García, C., Molowny, A., Garcia-Verdugo, J. M., & Ferrer, I. (1988). Delayed Postnatal Neurogenesis in the Cerebral-Cortex of Lizards. *Developmental Brain Research*, 43(2), 167-174.
- Marchioro, M., Nunes, J. M. D. M., Ramalho, A. M. R., Molowny, A., Perez-Martinez, E., Ponsoda, X., et al. (2005). Postnatal neurogenesis in the medial cortex of the tropical lizard *Tropidurus hispidus*. *Neuroscience*, 134(2), 407-413.
- Maurange, C., Cheng, L., & Gould, A. P. (2008). Temporal transcription factors and their targets schedule the end of neural proliferation in *Drosophila*. *Cell*, 133(5), 891-902.
- Mercier, F., Kitasako, J. T., & Hatton, G. I. (2002). Anatomy of the brain neurogenic zones revisited: fractones and the fibroblast/macrophage network. *The Journal of comparative neurology*, 451(2), 170-188.
- Meyer, R. L. (1978). Evidence from Thymidine Labeling for Continuing Growth of Retina and Tectum in Juvenile Goldfish. *Experimental Neurology*, 59(1), 99-111.
- Ming, G. L., & Song, H. (2005). Adult neurogenesis in the mammalian central nervous system. *Annu Rev Neurosci*, 28, 223-250.
- Molowny, A., Nacher, J., & Lopez-Garcia, C. (1995). Reactive neurogenesis during regeneration of the lesioned medial cerebral cortex of lizards. *Neuroscience*, 68(3), 823-836.
- Nowakowski, R. S., & Hayes, N. L. (2008). Numerology of Neurogenesis : Characterizing the Cell Cycle of Neurostem Cells In F. H. Gage, G. Kempermann & H. Song (Eds.), *Adult Neurogenesis* (pp. 9-10). New York: Cold Spring Harbor Laboratory Press.
- Pérez-Sánchez, F., Molowny, A., Garcia-Verdugo, J. M., & Lopezgarcia, C. (1989). Postnatal Neurogenesis in the Nucleus Sphericus of the Lizard, *Podarcis-Hispanica*. *Neuroscience Letters*, 106(1-2), 71-75.
- Pouwels, E. (1978). Development of Cerebellum of Trout, *Salmo-Gairdneri* .1. Patterns of Cell-Migration. *Anatomy and Embryology*, 152(3), 291-308.
- Raucci, F., Di Fiore, M. M., Pinelli, C., D'Aniello, B., Luongo, L., Polese, G., et al. (2006). Proliferative activity in the frog brain: A PCNA-immunohistochemistry analysis. *Journal of Chemical Neuroanatomy*, 32(2-4), 127-142.
- Raymond, P. A., Easter, S. S., Burnham, J. A., & Powers, M. K. (1983). Post-Embryonic Growth of the Optic Tectum in Goldfish .2. Modulation of Cell-Proliferation by Retinal Fiber Input. *Journal of Neuroscience*, 3(5), 1092-1099.
- Rowe, R. W. D., & Goldspink, G. (1969). Muscle fibre growth in five different muscles in both sexes of mice. *Journal of anatomy*(104, ), 519-530.
- Scharff, C., Kirn, J. R., Grossman, M., Macklis, J. D., & Nottebohm, F. (2000). Targeted neuronal death affects neuronal replacement and vocal behavior in adult songbirds. *Neuron*, 25(2), 481-492.

- Seri, B., Garcia-Verdugo, J. M., McEwen, B. S., & Alvarez-Buylla, A. (2001). Astrocytes give rise to new neurons in the adult mammalian hippocampus. *Journal of neuroscience*, 21(18), 7153-7160.
- Smart, I. (1961). Subependymal Layer of Mouse Brain and Its Cell Production as Shown by Radioautography after Thymidine-H3 Injection. *Journal of Comparative Neurology*, 116(3), 325-&.
- Song, H. J., Stevens, C. F., & Gage, F. H. (2002). Neural stem cells from adult hippocampus develop essential properties of functional CNS neurons. *Nature neuroscience*, 5(5), 438-445.
- Soutschek, J., & Zupanc, G. K. (1996). Apoptosis in the cerebellum of adult teleost fish, *Apteronotus leptorhynchus*. *Brain research. Developmental brain research*, 97(2), 279-286.
- Stanfield, B. B., & Trice, J. E. (1988). Evidence that granule cells generated in the dentate gyrus of adult rats extend axonal projections. *Experimental brain research.*, 72(2), 399-406.
- Temple, S. (2001). The development of neural stem cells. *Nature*, 414(6859), 112-117.
- Ueki, T., Tanaka, M., Yamashita, K., Mikawa, S., Qiu, Z., Maragakis, N. J., et al. (2003). A novel secretory factor, Neurogenesis-1, provides neurogenic environmental cues for neural stem cells in the adult hippocampus. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 23(37), 11732-11740.
- Zakon, H. H. (1984). Postembryonic changes in the peripheral electrosensory system of a weakly electric fish: addition of receptor organs with age. *The Journal of comparative neurology*, 228(4), 557-570.
- Zikopoulos, B., Kentouri, M., & Dermon, C. R. (2000). Proliferation zones in the adult brain of a sequential hermaphrodite teleost species (*Sparus aurata*). *Brain Behavior and Evolution*, 56(6), 310-322.
- Zupanc, G. K. H. (2008). Adult neurogenesis and neuronal regeneration in the brain of teleost fish. *Journal of Physiology-Paris*, 102(4-6), 357-373.
- Zupanc, G. K. H., Hinsch, K., & Gage, F. H. (2005). Proliferation, migration, neuronal differentiation, and long-term survival of new cells in the adult zebrafish brain. *Journal of Comparative Neurology*, 488(3), 290-319.
- Zupanc, G. K. H., & Horschke, I. (1995). Proliferation Zones in the Brain of Adult Gymnotiform Fish - a Quantitative Mapping Study. *Journal of Comparative Neurology*, 353(2), 213-233.
- Rowlerson, A., Veggetti, A., (2001). "Cellular mechanisms of post-embryonic muscle growth in aquaculture species". In: Johnston, I.A. (Ed.), *Muscle Development and Growth*. Academic Press, San Diego, pp. 103–140