

# **Proof Delivery Form**

Please return this form with your proof

**Article number: S0025315411000889jra**

**Date of delivery: 08.06.11**

**Typesetter ref number: MBI1100088**

**Volume and Issue Number: 0 and 0**

**Number of pages (not including this page): 10**

---

## **Journal of the Marine Biological Association of the United Kingdom**

Here is a pdf proof of your article for publication in the **Journal of the Marine Biological Association of the United Kingdom**. Please print out the file, check the proofs carefully and answer any queries.

Please return your corrections **via email** (no later than 4 days after receipt) quoting paper number in the header of the email message.

Please also ensure you specify page and line number of each correction required in your email and send to:

Executive Editor JMBA

Email: [jmba@mba.eclipse.co.uk](mailto:jmba@mba.eclipse.co.uk)

Please return your completed and signed **copyright transfer form** and **offprint form** by post to the addresses given on each form.

---

Please note:

- You are responsible for correcting your proofs. Errors not found may appear in the published journal.
- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal.
- Please answer carefully any queries raised from the typesetter.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required

Thank you in advance.

---

**Author queries:**

---

**Typesetter queries:**

---

Please return this form with your proof

# Offprint order form



**CAMBRIDGE**  
UNIVERSITY PRESS

PLEASE COMPLETE AND RETURN THIS FORM. WE WILL BE UNABLE TO SEND OFFPRINTS UNLESS A RETURN ADDRESS AND ARTICLE DETAILS ARE PROVIDED.

VAT REG NO. GB 823 8476 09

## Journal of the Marine Biological Association (MBI)

Volume:

no:

### Offprints

Offprints may be purchased at the prices shown below. Please complete this form and send it to **the publisher (address below)**. Please give the address to which your offprints should be sent. They will be despatched by surface mail within one month of publication.

Number of offprints required :

Email:

Offprints to be sent to (print in BLOCK CAPITALS):

Post/Zip Code:

Telephone:

Date (dd/mm/yy):

Author(s):

Article Title:

All enquiries about offprints should be addressed to **the publisher: Journals Production Department, Cambridge University Press, The Edinburgh Building, Shaftesbury Road, Cambridge CB2 8RU, UK.**

**Charges for extra offprints (excluding VAT) Please circle the appropriate charge:**

Number of copies	25	50	100	150	200	per 50 extra
1-4 pages	£68	£109	£174	£239	£309	£68
5-8 pages	£109	£163	£239	£321	£399	£109
9-16 pages	£120	£181	£285	£381	£494	£120
17-24 pages	£131	£201	£331	£451	£599	£131
Each Additional 1-8 pages	£20	£31	£50	£70	£104	£20

### Methods of payment

If you live in Belgium, France, Germany, Ireland, Italy, Portugal, Spain or Sweden and are not registered for VAT we are required to charge VAT at the rate applicable in your country of residence. If you live in any other country in the EU and are not registered for VAT you will be charged VAT at the UK rate.

If registered, please quote your VAT number, or the VAT number of any agency paying on your behalf if it is registered.

VAT Number:

Payment **must** be included with your order, please tick which method you are using:

- ☐ Cheques should be made out to Cambridge University Press.
- ☐ Payment by someone else. Please enclose the official order when returning this form and ensure that when the order is sent it mentions the name of the journal and the article title.
- ☐ Payment may be made by any credit card bearing the Interbank Symbol.

Card Number:

Expiry Date (mm/yy):

/

Card Verification Number:

The card verification number is a 3 digit number printed on the **back** of your **Visa** or **Master card**, it appears after and to the right of your card number. For **American Express** the verification number is 4 digits, and printed on the **front** of your card, after and to the right of your card number.

Signature of  
card holder:

Amount  
(Including VAT  
if appropriate):

£

Please advise if address registered with card company is different from above



## Transfer of copyright

Please read the notes overleaf and then complete, sign, and return this form to **The Executive Editor, Marine Biological Association of the UK, The Laboratory, Citadel Hill, Plymouth PL1 2PB, UK** as soon as possible.

In consideration of the publication in **JOURNAL OF THE MARINE BIOLOGICAL ASSOCIATION OF THE UNITED KINGDOM**

of the contribution entitled: .....

.....

by (all authors' names): .....

.....

### 1 To be filled in if copyright belongs to you

#### Transfer of copyright

I/we hereby assign to Cambridge University Press, full copyright in all formats and media in the said contribution, including in any supplementary materials that I/we may author in support of the online version.

I/we warrant that I am/we are the sole owner or co-owners of the contribution and have full power to make this agreement, and that the contribution contains nothing that is in any way an infringement of any existing copyright or licence, or duty of confidentiality, or duty to respect privacy, or any other right of any person or party whatsoever and contains nothing libellous or unlawful; and that all statements purporting to be facts are true and that any recipe, formula, instruction or equivalent published in the Journal will not, if followed accurately, cause any injury or damage to the user.

I/we further warrant that permission has been obtained from the copyright holder for any material not in my/our copyright including any audio and video material, that the appropriate acknowledgement has been made to the original source, and that in the case of audio or video material appropriate releases have been obtained from persons whose voices or likenesses are represented therein. I/we attach copies of all permission and release correspondence.

I/we hereby assert my/our moral rights in accordance with the UK Copyrights Designs and Patents Act (1988).

Signed (tick one)

☐ the sole author(s)    ☐ one author authorised to execute this transfer on behalf of all the authors of the above article

Name (block letters) .....

Institution/Company .....

Signature: ..... Date: .....

*(Additional authors should provide this information on a separate sheet.)*

### 2 To be filled in if copyright does not belong to you

a Name and address of copyright holder .....

.....

.....

b The copyright holder hereby grants to Cambridge University Press the non-exclusive right to publish the contribution in the journal and to deal with requests from third parties in the manner specified in paragraphs 4 and 5 overleaf.

(Signature of copyright holder or authorised agent) .....

### 3 US Government exemption

I/we certify that the paper above was written in the course of employment by the United States Government so that no copyright exists.

Signature: ..... Name (Block letters): .....

### 4 Requests received by Cambridge University Press for permission to reprint this article (see para. 4 overleaf) should be sent to

Name and address (block letters) .....

---

## Notes for contributors

- 1 The Journal's policy is to acquire copyright in all contributions. There are two reasons for this: (a) ownership of copyright by one central organisation tends to ensure maximum international protection against unauthorised use; (b) it also ensures that requests by third parties to reprint or reproduce a contribution, or part of it, are handled efficiently and in accordance with a general policy that is sensitive both to any relevant changes in international copyright legislation and to the general desirability of encouraging the dissemination of knowledge.
- 2 Two 'moral rights' were conferred on authors by the UK Copyright Act in 1988. In the UK an author's 'right of paternity', the right to be properly credited whenever the work is published (or performed or broadcast), requires that this right is asserted in writing.
- 3 Notwithstanding the assignment of copyright in their contribution, all contributors retain the following **non-transferable** rights:
  - The right to post *either* their own version of their contribution as submitted to the journal (prior to revision arising from peer review and prior to editorial input by Cambridge University Press) *or* their own final version of their contribution as accepted for publication (subsequent to revision arising from peer review but still prior to editorial input by Cambridge University Press) on their **personal or departmental web page**, or in the **Institutional Repository** of the institution in which they worked at the time the paper was first submitted, or (for appropriate journals) in PubMedCentral or UK PubMedCentral, provided the posting is accompanied by a prominent statement that the paper has been accepted for publication and will appear in a revised form, subsequent to peer review and/or editorial input by Cambridge University Press, in **Journal of the Marine Biological Association of the United Kingdom** published by Cambridge University Press, together with a copyright notice in the name of the copyright holder (Cambridge University Press or the sponsoring Society, as appropriate). On publication the full bibliographical details of the paper (volume: issue number (date), page numbers) must be inserted after the journal title, along with a link to the Cambridge website address for the journal. Inclusion of this version of the paper in Institutional Repositories outside of the institution in which the contributor worked at the time the paper was first submitted will be subject to the additional permission of Cambridge University Press (not to be unreasonably withheld).
  - The right to post the definitive version of the contribution as published at Cambridge Journals Online (in PDF or HTML form) on their **personal or departmental web page**, no sooner than upon its appearance at Cambridge Journals Online, subject to file availability and provided the posting includes a prominent statement of the full bibliographical details, a copyright notice in the name of the copyright holder (Cambridge University Press or the sponsoring Society, as appropriate), and a link to the online edition of the journal at Cambridge Journals Online.
  - The right to post the definitive version of the contribution as published at Cambridge Journals Online (in PDF or HTML form) in the **Institutional Repository** of the institution in which they worked at the time the paper was first submitted, or (for appropriate journals) in PubMedCentral or UK PubMedCentral, no sooner than **one year** after first publication of the paper in the journal, subject to file availability and provided the posting includes a prominent statement of the full bibliographical details, a copyright notice in the name of the copyright holder (Cambridge University Press or the sponsoring Society, as appropriate), and a link to the online edition of the journal at Cambridge Journals Online. Inclusion of this definitive version after one year in Institutional Repositories outside of the institution in which the contributor worked at the time the paper was first submitted will be subject to the additional permission of Cambridge University Press (not to be unreasonably withheld).
  - The right to post an abstract of the contribution (for appropriate journals) on the **Social Science Research Network (SSRN)**, provided the abstract is accompanied by a prominent statement that the full contribution appears in **Journal of the Marine Biological Association of the United Kingdom** published by Cambridge University Press, together with full bibliographical details, a copyright notice in the name of the journal's copyright holder (Cambridge University Press or the sponsoring Society, as appropriate), and a link to the online edition of the journal at Cambridge Journals Online.
  - The right to make hard copies of the contribution or an adapted version for their own purposes, including the right to make multiple copies for course use by their students, provided no sale is involved.
  - The right to reproduce the paper or an adapted version of it in any volume of which they are editor or author. Permission will automatically be given to the publisher of such a volume, subject to normal acknowledgement.
- 4 We shall use our best endeavours to ensure that any direct request we receive to reproduce your contribution, or a substantial part of it, in another publication (which may be an electronic publication) is approved by you before permission is given.
- 5 Cambridge University Press co-operates in various licensing schemes that allow material to be photocopied within agreed restraints (e.g. the CCC in the USA and the CLA in the UK). Any proceeds received from such licences, together with any proceeds from sales of subsidiary rights in the Journal, directly support its continuing publication.
- 6 It is understood that in some cases copyright will be held by the contributor's employer. If so, Cambridge University Press requires non-exclusive permission to deal with requests from third parties, on the understanding that any requests it receives from third parties will be handled in accordance with paragraphs 4 and 5 above (note that your approval and not that of your employer will be sought for the proposed use).
- 7 Permission to include material not in your copyright  
If your contribution includes textual or illustrative material not in your copyright and not covered by fair use / fair dealing, permission must be obtained from the relevant copyright owner (usually the publisher or via the publisher) for the non-exclusive right to reproduce the material worldwide in all forms and media, including electronic publication. The relevant permission correspondence should be attached to this form.

If you are in doubt about whether or not permission is required, please consult the Permissions Manager, Cambridge University Press, The Edinburgh Building, Shaftesbury Road, Cambridge CB2 8RU, UK. Fax: +44 (0)1223 315052. Email: [lnicol@cambridge.org](mailto:lnicol@cambridge.org).

The information provided on this form will be held in perpetuity for record purposes. The name(s) and address(es) of the author(s) of the contribution may be reproduced in the journal and provided to print and online indexing and abstracting services and bibliographic databases.

*Please make a duplicate of this form for your own records*

# Social structure of the declining resident community of common bottlenose dolphins in the Sado Estuary, Portugal

JOANA F. AUGUSTO<sup>1,2</sup>, PATRÍCIA RACHINAS-LOPES<sup>1,2</sup> AND MANUEL E. DOS SANTOS<sup>1,2</sup>

<sup>1</sup>Unidade de Investigação em Eco-Etologia, ISPA—Instituto Universitário, Rua Jardim do Tabaco 34, 1149-041 Lisboa, Portugal,

<sup>2</sup>Projecto Delfim—Centro Português de Estudo dos Mamíferos Marinhos, Rua Alto do Duque 45, 1400-009 Lisboa, Portugal

*The resident population of common bottlenose dolphins (*Tursiops truncatus*) in the Sado Estuary, Portugal, has been declining at least during the past three decades. A complete photographic census produced a current count of 24 animals—19 adults, three subadults and two calves. It appears to be phylopatric and essentially closed, but given the likely importance that exchanges with neighbouring coastal groups may play, even if rare, the most adequate term to define this dolphin should be community and not population. Large groups with all age-classes are common in the community, possibly as a calf and subadult protection strategy, and this may be related to the fact that these age-classes have had high mortality rates in the last decade. Maternity of two calves was determined, and we found that the two mothers adopted different parenting strategies. While one mother spent more time alone with her calf, the other mother spent more time with her calf in larger groups.*

*The average coefficient of association for this community is 0.45, quite high for this species. Associations and typical group size are similar between all individuals, with no patterning according to age-class or sex, which constitutes an atypical trait for dolphin societies. There are also no clear divisions in this community according to cluster analysis.*

*Associations are preferred and long term, lasting approximately 34 days and fitting a pattern of casual acquaintances, where individuals associate for a period of time, disassociate and may reassociate after that. This reflects the fission–fusion character of the community, but in a more stable manner. We think this is caused by a combination of demographic characteristics and a stable and productive environment, which led to a decrease in competition between individuals.*

**Keywords:** social structure, bottlenose dolphins, *Tursiops truncatus*, demographic effects, fission–fusion dynamics

Submitted 11 February 2010; accepted 16 May 2011

## INTRODUCTION

The social ecology and population biology of the common bottlenose dolphin, *Tursiops truncatus* (Montagu, 1821) have been studied in many locations around the world (reviewed by Connor *et al.*, 2000). It is a cosmopolitan species that occurs in temperate and tropical regions, both in coastal and deep waters (Reynolds *et al.*, 2000). Longevity in this species is up to 40 or 50 years, in males and females respectively. Females reach reproductive maturity between five and 13 years of age and males between eight and 13. The gestation period lasts 12 months, resulting in a single calf, with inter-birth intervals typically of three to four years (Connor *et al.*, 2000).

Throughout their extensive distribution range, common bottlenose dolphins are usually found in groups between two and 15 animals, although groups of hundreds or thousands have been reported in offshore waters (Scott & Chivers, 1990; Wells & Scott, 1994). Coastal populations form smaller groups, with a fission–fusion, dynamic composition, even though some associations may be very stable over the years (Connor *et al.*, 2000). Sex, age, reproductive

condition, familial relationships, affiliation histories or the formation of coalitions may influence the association patterns of these animals (Connor *et al.*, 1992; Wells & Scott, 1994; Connor & Whitehead, 2005; Whitehead & Connor, 2005).

Common bottlenose dolphins in the Sado Estuary region (central continental Portugal) form one of the few resident communities in Europe (Gaspar, 2003). The first reports of repeatedly recognizable individuals were presented in the early 1980s (Teixeira & Duguy, 1981) and later surveys indicated a clear residency pattern (dos Santos & Lacerda, 1987). Observations of interactions or exchanges between the resident group and adjacent, sympatric groups have always been rare or inconclusive (Gaspar, 2003).

Popular and anecdotal sources speak of over 50 dolphins in the area before the 1980s (dos Santos & Lacerda, 1987) and, using photo-identification, these authors estimated a local population of at least 40 individuals. Gaspar (2003) studied animals identified between 1981 and 1997 and listed 37 individuals considered resident at the end of this period. Survival rates varied according to age-class, with lower rates in calves and subadults, and these quantitative analyses supported the notion of rapid decline in this resident community.

Social structure may influence conservation, especially in species where strong bonds exist and local traditions affect movement or mating patterns (Sutherland, 1998;

**Corresponding author:**  
J.F. Augusto  
Email: jaugusto@ispa.pt

Whitehead, 2008a). Therefore, any insight into the association patterns within this declining community should be relevant to the evaluation of its survival.

Although this species is listed as 'Least Concern' in the 2009 update of the *IUCN Red List* (IUCN, 2009), local isolated populations of bottlenose dolphins may warrant the status of 'Critically Endangered' (Currey *et al.*, 2009). Recognizing the grave condition of the local population in the Sado region, the conservation authority in Portugal, Instituto da Conservação da Natureza e Biodiversidade, has approved an Action Plan to protect and monitor these bottlenose dolphins (Sequeira *et al.*, 2009).

As stated in the Action Plan, while demographic and behavioural information on the bottlenose dolphin community resident in the Sado Estuary is limited, the assessment of its conservation status will benefit from the analysis of its social structure in the greatest detail possible.

With this study we analyse the social structure of this community, relating our results to the available demographic data and considering them from a conservation perspective.

## MATERIALS AND METHODS

### Study area

The bottlenose dolphin groups were observed and sampled in the core region of this population's home range: the Sado Estuary and the adjacent coastal waters (Figure 1), where these animals feed and socialize on a year-round basis. The study area covers about 200 km<sup>2</sup> and is rather variable in physiography and level of human influence: there are a city front and a busy harbour; long and quiet beaches and rocky shores; shipping lanes and also protected areas. The river mouth, 40 m deep, is located at approximately 38°29'28"N 08°55'28"W. Industrial pollutants, herbicides and pesticides have been accumulating for decades in the water and

sediments, reaching high contamination levels in some areas (e.g. Gil & Vale, 2001; Caeiro *et al.*, 2005).

### Data collection

Group composition data, photographic and behavioural records were collected during 40 days, from April 2007 to July 2010, a total of 195 hours, with at least three observers on board an 8.40-m motor launch. Groups were followed focally from a distance between 50 and 100 m. Using the definition of Shane (1990a, b), groups are sets of individuals in apparent association, moving in the same direction, usually engaged in the same behaviour. Photographic records of the individuals were collected with a digital Nikon D70S or/and a Canon EOS 400D (both with 70–300 mm zoom lenses).

### Data analysis

Photographs were used to identify individuals from the marks and scars on their dorsal fins (Würsig & Würsig, 1977), and these records were compared to the accumulated photo catalogue started in 1981 and managed jointly by ISPA—Instituto Universitário and Projecto Delfim. Age-class and sex for each individual identified were determined using previous data (e.g. dos Santos & Lacerda, 1987; Harzen & dos Santos, 1992; Gaspar, 2003).

The method described in Grellier *et al.* (2003) was used to determine maternity. In this method, photographic records from the calf's first sighting until the last sighting the next calendar year were used to calculate coefficients of association (CoAs) using the simple ratio index ( $p$ )

$$p = \frac{x}{y_a + y_b}$$

where  $x$  is the number of times a calf (individual A) was seen with another identified animal (individual B),  $y_a$  the number




Fig. 1. Map of the study area, on the central western coast of Portugal. The small dotted lines represent mud banks and the broken lines represent Marine Protected Areas and RNES (Natural Reserve of the Sado Estuary).



of times where the calf was sighted without individual B, and  $y_b$  the number of times where the individual B was sighted without the calf. The sum of  $y_a$  and  $y_b$  is noted as  $n$ , the total number of times either animal was seen. The standard error of  $p$  is estimated as

$$SE = \sqrt{\frac{p(1-p)}{n}}$$

A one tailed z-test was then used to compare the CoAs of the top associates of the calf

$$z = \frac{p_1 - p_2}{\sqrt{p(1-p)(\frac{1}{n_1} + \frac{1}{n_2})}}$$

This is only considered a good approach when  $n_1 + n_2 > 12$  (Grellier *et al.*, 2003).

The CoAs are calculated using two different approaches, members of the same group (defined as in Shane, 1990a, b) and in the same photographic frame. Analysis was first carried out using groups and, if according to the z-test, the CoA of the top associate was significantly greater than the other associate, the top associate was designated as the calf's mother. If the z-test determined that both top associates had a similar CoA, the analysis was repeated using the photographic frames. If according to the z-test, the CoA of the top associate was not significantly greater than the other associate in both analyses the maternity of the calf remained undetermined.

Photographs allowed the confirmation of estimated group sizes and group composition based on visual counts and identifications. Only data confirmed by photographic evidence were used. Groups were categorized according to the age-classes present (procedure adapted from Félix, 1997): big and robust individuals, most of them previously identified and catalogued, were considered adults; less robust and smaller individuals, not in close association with a particular adult, were considered subadults; small individuals, sometimes with fetal folds, in close association with an individual were considered calves. Average group sizes according to class were analysed with a Kruskal-Wallis test (Zar, 1996) using Statistica v. 8.0 (StatSoft, Inc.). The null hypothesis is that average group sizes are similar between groups with different age-classes.

Typical group size (see Jarman, 1974, appendix 2), the size as experienced by an individual, was also estimated

$$tgs = \frac{\sum N_g(i)^2}{\sum N_g(i)}$$

where  $N_g$  is the count of the size  $n$  of observed groups.

Coefficients of association between dyads were calculated with SOCPROG 2.3 (Whitehead, 2009) using the half weight index (HWI), considered less biased for this kind of sampling (Cairns & Schwager, 1987):

$$HWI = \frac{x}{x + 1/2(y_a + y_b)}$$

in which  $x$  is the number of groups where the individuals A and B were seen together,  $y_a$  the number of groups where

the individual A was sighted without the individual B, and  $y_b$  the number of groups where the individual B was sighted without the individual A.

Coefficients vary between 0 (individuals never seen together) and 1 (individuals always seen together). Only individuals identified in more than 20 photographs were used. The standard error associated with each CoA was calculated using the binomial approach:

$$SE = \sqrt{\frac{a(1-a)}{n}}$$

in which  $a$  is the calculated CoA and  $n$  is the total number of times either animal was seen.

The randomness of CoAs was tested using the preferred/avoided associations test (Bejder *et al.*, 1998; Whitehead, 1999; Whitehead *et al.*, 2005). The null hypothesis for this test is that associations between pairs of individuals are random, and it is refuted if the standard deviation of the calculated coefficients is significantly higher than those from the permuted data. The routine chosen is SOCPROG 2.3 tests for both short and long term associations. If the standard deviation of calculated associations is significantly higher than the permuted data, the associations are long term preferred; if the mean association of calculated CoAs is significantly lower than the permuted, the associations are short term preferred.

Gregariousness, or differences in sociality between individuals (Whitehead *et al.*, 2005) was tested, in order to look for individuals that may be consistently found in groups larger or smaller than the typical (*sensu* Jarman, 1974), also using data from the permutation tests. The null hypothesis is that all individuals are found in groups with a similar distribution of sizes. It is refuted when the standard deviation of typical group size (SDtgs) has unexpectedly high values that are significantly different to the permuted data.

Accuracy of the social representation was assessed using  $r$ , the correlation coefficient between the estimated CoAs and the true CoAs (the proportion of time a pair is actually associated) (Whitehead, 2008b).

$$r = \frac{S}{CV(\alpha_{AB})}$$

in which  $S$  is the social differentiation, the estimated CV of the true CoAs and  $CV(\alpha_{AB})$  the CV of estimated CoAs. Values vary between 0, not an accurate representation, to 1, an excellent representation.

$S$  indicates how variable the CoAs are within a population. Values smaller than 0.3 indicate CoAs between individuals are similar, so relationships within the population are homogeneous,  $S$  greater than 0.5 indicates that relationships are well varied and a  $S$  greater than 1 represents very varied and differentiated relationships (Whitehead, 2008b).

Social organization of the community was graphically displayed in a dendrogram using average linkage hierarchical cluster analysis (Morgan *et al.*, 1976; Colgan, 1978). To assess the level of division of the population into communities the modularity of the clustering was calculated (Newman, 2004, 2006). Modularity, modified for weighted networks and applied to coefficients of association, can be described as the difference between the expected proportion of

association within clusters—in this case, sets of individuals—and the proportion obtained with the data (Newman, 2006). A modularity different from zero indicates that there is a deviation from random, but only values superior to 0.3 are considered good indicators of division. The maximum modularity possible is 1, when members of different clusters do not associate. For these analyses only individuals that are currently alive and are not calves were taken into account, since the low CoAs from deceased individuals are not caused by avoidance of others, and calves associate closely with their mothers during their first years, thus they do not truly reflect the association patterns.

To model how associations vary with time a standardized lagged association rate (SLAR) analysis was performed (Whitehead, 1995) using all available data. This rate is the probability that, if two individuals, A and B, are associated at a particular time, then  $\tau$  units of time later, a randomly chosen association of individual A will be B. The sampling period used was 1 hour, and the SLAR was compared to the null association rate, i.e. the SLAR if associations were random.

The SLAR obtained was also compared with theoretical models of different types of social structure (Whitehead, 1995). To assess which was the one most similar to our data the quasi-Akaike information criterion (QAIC) was calculated. The model that minimized this criterion was considered the best fit (Whitehead, 2007). The fit of the other models was also assessed using the variation of QAIC ( $\Delta$ QAIC). If  $\Delta$ QAIC is between 0 and 2 there is substantial support for the model, if it is between 4 and 7 it has considerably less support and if it is larger than 10 it has no support (Burnham & Anderson, 2002).

## RESULTS

### Community composition and demographic structure

A complete census revealed that the Sado resident community is currently composed of 24 individuals. A total of 13,035 photographs, collected from April 2007 to July 2010 (Table 1), were analysed, 63% of which were used for photo-id purposes. From these, 11,303 identifications of 27 individuals were obtained (Table 2). However, during the sampling period three of the individuals disappeared (LIN,

RED and TUD). All individuals were identified on the first sampling day, except the calves born in the summer of 2007—HUX and LIN—;summer of 2010—TAI; and an individual—TIP—who returned to the community in June 2007 after an apparent absence of at least one year.

Currently, this community is comprises 79.2% adults, 12.5% subadults and 8.3% calves, so one of its most striking features is the vast majority of adults. Half of the current community (and 63.1% of adults) have been observed since the 1980s, their first sighting varying from 1981 to 1985. These individuals were already adults at the time of first sighting, so they should be at least currently 30 years old. The sex of most individuals is unknown (58.3%), and most of those identified are females (29.2%).

### Testing for maternity of calves

According to the maternity test, GOR is DAR's mother (Table 3) and AGU is HUX's mother (Table 3). It was not possible to pinpoint LIN's mother because of the calf's low number of identifications and the analysis for TAI could not be carried out because the test is only performed with photographic records spanning a period of over one year.

### Group size and associations

A total of 258 groups was analysed, their size varying between one and 26 individuals. Average group size was  $7.75 \pm 6.37$  (Table 4), i.e. about 1/3 of the community size. Typical group size for this community is 12.97, i.e. just over half of

**Table 2.** Photoidentifications for each individual identified in the population, from April 2007 to July 2010. ID, identification of individual; Nr. IDs, number of photographs where the individual was identified in; Nr. Groups, number of groups where the individual was identified in.

ID	Nr. IDs	Nr. Groups
AGU	608	101
APA	495	88
BUM	670	104
CAL	384	74
CLU	465	103
DAR	479	89
ELE	304	59
FAC	523	106
GOR	549	99
HUX	377	73
LAM	685	112
LIN	8	6
LUA	585	107
MED	395	66
MID	617	80
MUR	660	116
QUA	520	74
RED	174	23
SPI	349	65
TAI	38	13
TAL	308	54
THO	507	89
TIP	244	39
TRU	535	90
TUD	158	33
WAL	352	68
ZOE	268	59

**Table 1.** Monthly distribution of sampling effort from 2007 to 2010.

	2007	2008	2009	2010	Total
January					
February		1		1	2
March		1	1		2
April	1			1	2
May	1				1
June	1	1	3	2	7
July	6		3	4	13
August	1	2	3		6
September		1			1
October	1				1
November					
December					
Total	11	6	10	8	35



**Table 3.** Maternity test according to Grellier *et al.* (2003). Significant test results are marked in bold ( $z_{0.05} = 1.64$ ). F. 1 High., female with the highest coefficient of association (CoA) with the calf; F. 2 High., female with the second highest CoA with the calf; p, CoA using simple ratio; n1, total number of times F.1 High and calf were seen together; n2, total number of times F.2 High and calf were seen together; z, unicaudal z-test result.

Calf	F.1 High.	p	F.2 High.	p	n1 + n2	z	Sampling unit
DAR	GOR	0.70	ELE	0.46	128	<b>2.75</b>	Group
HUX	RED	0.56	AGU	0.52	61	0.31	Frame
HUX	AGU	0.14	TRU	0.05	521	<b>3.41</b>	

the community size. The most frequent group categories were 'All adults' and 'Adults, subadults and calves' while the less frequent were 'Subadults and calves' and 'Calves' (Table 4). No groups with 'Subadults only' were recorded. According to the Shapiro–Wilk normality test ( $P < 0.001$ ), the size distribution according to classes is not normal, so a non-parametric approach was used, through a Kruskal–Wallis test. Average group size varies according to class (Table 5): groups with adults, subadults and calves have the largest average size in this community.

The association matrix resulting from the HWI (see Appendix) has an average of  $0.45 \pm 0.15$ . Most of the CoAs are medium (0.41–0.60) and there is only one high CoA ( $> 0.81$ ) (Figure 2). The association patterns are displayed in a dendrogram (Figure 3), which represents the data accurately, with a cophenetic correlation coefficient of 0.83.

SOCPROG estimated a correlation value of 0.942 between the CoAs calculated using our sampled data and the true associations, so the calculated CoAs are a good representative of the real ones and conclusions drawn from them can be applied to the community. The permutation tests determined that associations between dyads are preferred and long term, since the standard deviation of calculated CoAs is significantly higher than the permuted dataset (Table 6). There are no differences in gregariousness in this community, since the SDtgs is low and has a corresponding low value when compared to the permuted data (Table 6). This community has a homogeneous structure, with a social differentiation value of 0.38. Cluster analysis does not show a clear division in the community, given the maximum modularity of 0.05.

The SLAR analysis shows that association rates vary trough time, but remain above the null rate until at least 1000 days (Figure 4). The large error bars at the seven day mark reflect the structure of the data. Since there are not many consecutive sampling days, how associations vary on the short term is not very well understood. The error bars become smaller because the longer the time frame, the more data were collected, hence our understanding of variation in associations increased. The

**Table 4.** Groups according to classes for data from 2007 to 2010. N, number of samples; %, percentage of group-class in the sample;  $\bar{X}$ , average group size; SD, standard deviation.

Group-class	N	%	$\bar{X}$	SD
All adults	82	31.78	3.04	2.61
All subadults	0	0.00		
Adults and subadults	74	28.68	6.26	3.81
Adults and calves	20	7.75	6.30	3.40
Adults, subadults and calves	79	30.62	14.63	5.91
Subadults and calves	2	0.78	2.00	0.00
Calves	1	0.39	1.00	2.61
Total	258		7.75	6.37

best fit model for the associations in this community is the one consisting of casual acquaintances (Table 7),  $g(t) = a \cdot e^{(-bt)} = 0.045345 \cdot e^{-(t \cdot 4.4061 \times 10^{-5})}$ . In this scenario, the duration of associations can be estimated by  $1/b$  (Whitehead, 2008a). Given that  $b = 4.4061 \times 10^{-5}$  associations last approximately 34 days in this community.

## DISCUSSION


A complete census of the Sado resident community of bottlenose dolphins has been conducted, with an analysis of the available demographic information and of social structure from April 2007 to July 2010.

This is not only a very small, but also an aged community, since 63.1% of adults are, at least, 30 years old. During the 1990s none of the calves born in this community survived for more than two years (Gaspar, 2003), causing the current imbalance between age-classes. Still, there is no evidence of reproductive senescence in this species, as females may be reproductively active until their late forties (Cockcroft & Ross, 1990; Wells & Scott, 1994; Reynolds *et al.*, 2000), so it is not unreasonable to expect them to be reproductively active after that age. This notion is supported by the fact that AGU was estimated to be at least 31 years old when HUX was born and that GOR was estimated to be at least 27 years old when DAR was born.

Since these dolphins have been studied there was only one documented case of an individual that was seen with an adjacent, non-resident group and then returned to the resident community (Gaspar, 2003). During our sampling period only one animal was apparently absent for at least one year, possibly joining other groups. However, no contacts between resident and non-resident animals have been observed in this study. Most likely this community is strongly phylopatric and has been essentially closed. However, considering the likely relevance of those exchanges, even if they are

**Table 5.** Kruskal–Wallis test results for average group size according to classes. Significant  $P$  value for 0.05 significance is marked in bold. N, number of samples;  $\bar{X}$ , average group size; S.R., sum of ranks; H, Kruskal–Wallis statistics;  $P$ ,  $P$  value.

Group-class	N	$\bar{X}$	S.R.
All adults	82	3.04	5346.50
All subadults	0		
Adults and subadults	74	6.26	9214.50
Adults and calves	20	6.30	2552.00
Adults, subadults and calves	79	14.63	16193.00
Subadults and calves	2	2.00	90.00
Calves	1	1.00	15.00
$H(5, N = 258) = 147.90$			<b><math>P &lt; 0.001</math></b>




**Fig. 2.** Coefficients of association classified according to Quintana-Rizzo & Wells (2001): low 0.01–0.20; medium–low 0.21–0.40; medium 0.41–0.60; medium–high 0.61–0.80; and high 0.81–1.

rare, the term community (referring to a group of animals in which most individuals interact with most others: Whitehead, 2008a, p. 14), instead of population, is perhaps the most adequate designation for the biosocial status of these dolphins.

Average group size is similar to previous studies in the Sado area (dos Santos & Lacerda, 1987; Harzen, 1995) and similar to the one recorded in previous studies elsewhere (Scott & Chivers, 1990; Wells & Scott, 1994), but the typical group size (as experience by the individuals) is much larger, just over half of the current community size. Large groups with all age-classes are quite common, possibly representing a calf and subadult protection strategy, as suggested by Kerr *et al.* (2005) and Gowans *et al.* (2008). Younger classes have had a high mortality rate in the last decade, and contamination, fishing nets and other forms of habitat degradation are all possible causes (Gaspar, 2003). The non-existence of ‘Subadults only’ groups in this study period may also be related to this protection strategy.

Interestingly, according to the results of the maternity tests, GOR and AGU adopted different strategies with their calves. Since the GOR could be determined as DAR’s mother using



**Fig. 3.** Dendrogram of the hierarchical cluster analysis using average linkage for all individuals currently alive in the population, except calves. The x axis represents the coefficients of association values, and the y axis represents the individuals.

**Table 6.** Preferred/avoided associations test—comparison between real and permuted coefficients of association (CoAs). Permuted data were calculated using 1000 random permutations and 10,000 trials per permutation. Significant *P* values for 0.05 significance are marked in bold.  $\bar{X}$ , average CoAs; SD, standard deviation; CV, coefficient of variation; SD tgs, standard deviation of typical group size.

	$\bar{X}$	SD	CV	SD tgs
Calculated	0.45	0.15	0.33	1.01
After permutation	0.18	0.05	0.31	1.15
<i>P</i> value	<b>0.001</b>	<b>&lt;0.001</b>	<b>0.001</b>	0.97

only associations based on group membership it means the pair spent a considerable amount of time separate from other groups, while the opposite happened with AGU. She could only be identified as HUX’s mother using data on photographic frames, so the pair was found consistently in groups with other individuals but in close spatial association.


It is unclear why these females adopted different strategies, but it is possible that patterns of association could change according to the calf’s age. We know that DAR was born in summer 2006, but data available for this analysis were only recorded since March 2007. Therefore, results for the maternity test might be reflecting the parenting strategy after the first six months of the calf’s life.

Bottlenose dolphin societies are characterized by a fission–fusion dynamic, with group membership varying within a very small time frame, leading to low association coefficients between pairs of individuals (with the exception of some long lasting bonds, usually between alliance forming males and mother–calf pairs) and consequently to a low average coefficient of association for the community.

The Sado community shows a different social organization. It presents an average coefficient of association of 0.45, higher than other resident communities, which vary between 0.1 and 0.3 (Smolker *et al.*, 1992; Félix, 1997; Connor *et al.*, 2000; Quintana-Rizzo & Wells, 2001; Chilvers & Corkeron, 2002; Eisefeld & Robinson, 2004). It is also higher than what was previously calculated for this resident community (Harzen, 1995), when its size was larger and most similar to Doubtful Sound’s population ( $0.47 \pm 0.04$ ) (Lusseau *et al.*, 2003). A small community size influences the associations only up to a point. A good estimator of the average coefficient of associations for a population is typical group size divided by population size minus 1 (Whitehead, 2008a). So, coefficients of association are affected not only by population size—the smaller the population, the larger the average coefficient—but also by typical group size. It is possible to have a small population size with a low average coefficient of association if the typical group size is also small. For example, if this population had a typical group size of 8 individuals (similar to average group size) the average coefficient of association would be closer to 0.3, which would be an average value for this species.

Given the social differentiation, relationships within this community are homogeneous, so pairs of individuals have similar associations. There are no differences in gregariousness, so all individuals have a similar typical group size. These are all atypical traits for bottlenose dolphin societies, in which association patterns are commonly influenced by factors such as the age and sex of the individuals.

Associations are also preferred and long term, so individuals associate in a non-random and preferential way, with associations lasting for approximately 34 days, according to



**Fig. 4.** Standardized lagged association rate (SLAR) for all individuals. Error bars were calculated using the jackknife technique. The null association rate represents the theoretical SLAR if individuals associated randomly. The maximum-likelihood best fit model represents casual acquaintances.

the SLAR. The patterns of association seem to be best described as casual acquaintances, individuals that associate for a period of time, dissociate after it and may reassociate afterwards. This obviously reflects the fission–fusion dynamic typical for these societies, although even more stable than previously reported (Harzen, 1995). Community size is known to influence fission–fusion societies in wild chimpanzees (Lehman & Boesch, 2004). Cohesiveness increases and fission–fusion patterns become less flexible with small communities. It is possible that community size has been influencing the Sado social structure in the same manner.

The only bottlenose dolphin population that, so far, has been found to have a stable social structure is resident in Doubtful Sound (Lusseau *et al.*, 2003). Other resident bottlenose dolphin populations have much more fluid social structures, as is reflected by their low average CoA (Smolker *et al.*, 1992; Félix, 1997; Connor *et al.*, 2000; Quintana-Rizzo & Wells, 2001; Chilvers & Corkeron, 2002; Eisfeld & Robinson, 2004). Although the Sado community does not show as long lasting associations, they still last longer than would be expected in a typical fission–fusion society and it is interesting to compare both populations. There are several parallelisms between the resident population in Doubtful Sound and in the Sado estuary. They are both small, nearly

closed and phylopatric, although Doubtful Sound still has a population of 65 individuals (Lusseau *et al.*, 2003). The main difference relates to the habitat type. While Doubtful Sound is a fjord with highly variable spatial and temporal productivity, the Sado Estuary provides stable feeding resources for the bottlenose dolphins throughout the year. With a decreasing number of individuals in such a stable feeding ground, it is likely that intraspecific competition has diminished, resulting in a high typical group size and increasing the stability of the association patterns (Perrin & Lehmann, 2001; Gowans *et al.*, 2008). The closed and phylopatric nature of this community, and its dependence on its habitat, affects its social structure but it also renders it more fragile, as it is unlikely to benefit from immigration or exchanges with other groups. If contact with other groups is happening, in order for it to not be detected in the present study, it must be sporadic or located in a time frame when sampling was not performed—nighttime or winter. Future work should focus on collecting data on coastal non-resident groups and their possible interactions with the resident community. This would make it possible to apply social network models and determine how the Sado community interacts with neighbouring groups. It would be interesting to assess if contacts are carried out by specific individuals—brokers (Lusseau & Newman, 2004)—instead of by the community as a whole and to model how contacts are affected by the loss of certain individuals. Different individuals may have different connectivity within or between communities and their removal may cause grave effects on the community (Lusseau & Newman, 2004). Contacts might have a particular importance for this community, especially at this low number of individuals.

In Little Bahama Bank, two hurricanes caused the loss of 30% of a population that comprised only one community (Elliser & Herzing, 2010). The remaining individuals formed two separate communities, each integrating immigrants. This population initially comprised 190 individuals, a much larger number than the Sado community. It also had a more fluid social structure and it was not geographically or demographically isolated from other populations, which may explain why immigrants were so well accepted.

It would be likely that if the Sado community size continues decreasing the fission–fusion dynamics will continue stabilizing, but how it would affect contacts with other populations is still unknown.

With this study we show how the social structure of the Sado community is affected by its demography. Its small

**Table 7.** Fit of social models to the standardized lagged association rate for the community. t, time in days; QAIC, quasi-Akaike information criterion;  $\Delta$ QAIC, variation of QAIC between the current model and the best fit; g, SLAR; td, time (days).

Model description	Model formula	Maximum-likelihood values for parameters	Standard errors for parameters (jackknife)	Number of parameters	QAIC	$\Delta$ QAIC
Constant companions (CC)	$g(t) = a$	$a = 0.044401$	$8.9318 \times 10^{-4}$	1	540211.8174	16.0095
Casual acquaintances (CA)	$g(t) = a.e^{(-bt)}$	$a = 0.045345$ $b = 4.4061 \times 10^{-5}$	$1.7374 \times 10^{-3}$ $4.2649 \times 10^{-5}$	2	540195.8079	
CC + CA	$g(t) = a + c.e^{(-bt)}$	$a = 0.044404$ $b = 1.5484$ $c = -8.3827 \times 10^{-3}$	$8.9031 \times 10^{-4}$ 17.4778 145.8831	3	540215.645	19.8371
Two levels of CA	$g(t) = a.e^{(-bt)} + c.e^{(-dt)}$	$a = 0.044052$ $b = 24.8201$ $c = 0.045347$ $d = 4.4132.10^{-5}$	14.5478 53.2079 $1.7326 \times 10^{-3}$ $4.2708 \times 10^{-5}$	4	540199.8079	4



and declining size, in a stable feeding ground, has led to a decrease in intraspecific competition and, consequently, to a more stable association pattern of fission–fusion dynamics. We believe that detailed information on the social structure of this community will promote a better understanding of its conservation perspectives which may depend on contacts with non-resident groups.

## ACKNOWLEDGEMENTS

We are grateful to all our colleagues who participated in the fieldwork and data collection, especially Miguel N. Couchinho, Ana Rita Luís, Cecília V. Ferreira, Sónia Louro, Ana Teresa Cândido, Erica Sá and Carina Silva. Thanks are due to Miguel N. Couchinho for the map of the study area, to Hal Whitehead for his comments on the final draft, and to the anonymous referees who helped to improve the manuscript. We also acknowledge with gratitude all the current supporters of Projecto Delfim, especially Zoomarine, Alcatel, Via Verde and Companhia das Sandes. This work was supported by the Portuguese Foundation for Science and Technology (FCT) (Grant number MAR-LVT-Lisboa-331), and by private sponsors (Zoomarine-Portugal and Alcatel).

## REFERENCES

- Bejder L., Fletcher D. and Bräger S. (1998) A method for testing association patterns of social animals. *Animal Behaviour* 56, 719–725.
- Burnham K.P. and Anderson D.R. (2002) *Model selection and multimodel inference: a practical information-theoretic approach*. 1st edition. New York: Springer-Verlag.
- Caeiro S., Costa M.H., Ramos T.B., Fernandes F., Silveira N., Coimbra A., Medeiros G. and Painho M. (2005) Assessing heavy metal contamination in Sado Estuary sediment: an index analysis approach. *Ecological Indicators* 5, 151–169.
- Cairns S.J. and Schwager S.J. (1987) A comparison of association indices. *Animal Behaviour* 35, 1454–1469.
- Chilvers B.L. and Corkeron P.J. (2002) Association patterns of bottlenose dolphins (*Tursiops aduncus*) off Point Lookout, Queensland, Australia. *Canadian Journal of Zoology* 80, 973–979.
- Cockcroft V.G. and Ross G.J.B. (1990) Age, growth, and reproduction of bottlenose dolphins *Tursiops truncatus* from the east coast of southern Africa. *Fishery Bulletin* 88, 289–302.
- Colgan P.W. (1978) Hierarchical cluster analysis. In Colgan P.W. (ed.) *Quantitative ethology*. 1st edition. New York: J. Wiley & Sons, pp.116–144.
- Connor R.C., Smolker R.A. and Richards A.F. (1992) Two levels of alliance formation among male bottlenose dolphins (*Tursiops* sp.). *Proceedings of National Academy of Sciences of the United States of America* 89, 987–990.
- Connor R., Wells R.S., Mann J. and Read A.J. (2000) The bottlenose dolphin: social relationships in a fission–fusion society. In Mann J., Connor R.C., Tyack P.L. and Whitehead H. (eds) *Cetacean societies: field studies of dolphins and whales*. 2nd edition, Chicago: University of Chicago Press, pp. 91–126.
- Connor R. and Whitehead H. (2005) Alliances II. Rates of encounter during resource utilization: a general model of intrasexual alliance formation in fission–fusion societies. *Animal Behaviour* 69, 127–132.
- Currey R.C.J., Dawson S.M. and Slooten E. (2009) An approach for regional threat assessment under IUCN Red List criteria that is robust to uncertainty: the Fiordland bottlenose dolphins are critically endangered. *Biological Conservation* 142, 1570–1579.
- dos Santos M.E. and Lacerda M. (1987) Preliminary observations of the bottlenose dolphin (*Tursiops truncatus*) in the Sado estuary (Portugal). *Aquatic Mammals* 13, 65–80.
- Eisfeld S.M. and Robinson K.P. (2004) The sociality of bottlenose dolphins in the outer southern Moray Firth, NE Scotland: implications for current management proposals? *European Research on Cetaceans* 18. [on CD-ROM.]
- Elliser C.R. and Herzog D.L. (2010) Replacement dolphins? Social restructuring of a resident pod of Atlantic bottlenose dolphins, *Tursiops truncatus*, after two major hurricanes. *Marine Mammal Science* 27, 39–59. Published online: 15 July 2010. DOI: 10.1111/j.1748-7692.2010.00403.x
- Félix F. (1997) Organization and social structure of the coastal bottlenose dolphin *Tursiops truncatus* in the Gulf de Guayaquil, Ecuador. *Aquatic Mammals* 23, 1–16.
- Gaspar R. (2003) *Status of the resident bottlenose dolphin population in the Sado estuary: past, present and future*. PhD thesis. St Andrews University, Scotland.
- Gil O. and Vale C. (2001) Evidence for polychlorinated biphenyls dechlorination in the sediments of Sado Estuary, Portugal. *Marine Pollution Bulletin* 42, 453–461.
- Gowans S., Würsig B. and Karczmarski L. (2008) The social structure and strategies of delphinids: predictions based on an ecological framework. *Advances in Marine Biology* 53, 195–294.
- Grellier K., Hammond P.S., Wilson B., Sanders-Reed C.A. and Thompson P.M. (2003) Use of photo-identification data to quantify mother–calf association patterns in bottlenose dolphins. *Canadian Journal of Zoology* 81, 1421–1427.
- Harzen S. (1995) *Behaviour and social ecology of the bottlenose dolphin, Tursiops truncatus, in the Sado estuary, Portugal*. PhD thesis. University of Bielefeld, Germany.
- Harzen S. and dos Santos M.E. (1992) Three encounters with bottlenose dolphins (*Tursiops truncatus*) carrying dead calves. *Aquatic Mammals* 18, 49–55.
- IUCN (2009) *2009 IUCN Red List of Threatened Species*. Version 2009.2. <<http://www.iucnredlist.org>> (accessed 2 February 2010).
- Jarman P.J. (1974) The social organization of antelope in relation to their ecology. *Behaviour* 48, 215–267.
- Kerr K.A., Defran R.H. and Campbell G.S. (2005) Bottlenose dolphins (*Tursiops truncatus*) in the Drowned Cayes, Belize: group size, site fidelity and abundance. *Caribbean Journal of Science* 41, 172–177.
- Lehman J. and Boesch C. (2004) To fission or to fusion: effects of community size on wild chimpanzee (*Pan troglodytes verus*) social organization. *Behavioral Ecology and Sociobiology* 56, 207–216.
- Lusseau D. and Newman M.E.J. (2004) Identifying the role that animals play in their social networks. *Proceedings of the Royal Society of London B (Supplement)* 271, 447–481.
- Lusseau D., Schneider K., Boisseau O.J., Haase P., Slooten E. and Dawson S.M. (2003) The bottlenose dolphin community of Doubtful Sound features a large proportion of long-lasting associations. Can geographic isolation explain this unique trait? *Behavioral Ecology and Sociobiology* 54, 396–405.
- Morgan B.J.T., Simpson M.J.A., Hanby J.P. and Hall-Craggs J. (1976) Visualizing interaction and sequential data in animal behaviour: theory and application of cluster-analysis methods. *Behaviour* 56, 1–43

- Newman M.E.J.** (2004) Analysis of weighted networks. *Physical Review E* 70, 056131 1–9.
- Newman M.E.J.** (2006) Modularity and community structure in networks. *Proceedings of the National Academy of Sciences of the United States of America* 103, 8577–8582.
- Perrin N. and Lehmann L.** (2001) Is sociality driven by the costs of dispersal or the benefits of phylopatriy? A role for kin-discrimination mechanisms. *American Naturalist* 158, 471–483.
- Quintana-Rizzo E. and Wells R.S.** (2001) Resighting and association patterns of bottlenose dolphins (*Tursiops truncatus*) in the Cedar Keys, Florida: insights into social organization. *Canadian Journal of Zoology* 79, 447–456.
- Reynolds III J.E., Wells R.S. and Eide S.D.** (2000) *The bottlenose dolphin. Biology and conservation*. 1st edition. Florida: University Press of Florida.
- Sequeira M., Matias S., Farinha J.C., Gaspar R., Silva C., Augusto J., Ferreira C.V., Fonseca M.J., Narra P. and Luís A.R.** (2009) *Bases para o Plano de Acção para a Salvaguarda e Monitorização da População de Roazes do Estuário do Sado*. 1st edition. Lisbon: Instituto da Conservação da Natureza e da Biodiversidade. [In Portuguese.]
- Scott M.D. and Chivers S.J.** (1990) Distribution and herd structure of bottlenose dolphins in the eastern tropical Pacific Ocean. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. 1st edition. San Diego, CA: Academic Press, pp. 387–402.
- Shane S.H.** (1990a) Behavior and ecology of the bottlenose dolphin at Sanibel Island, Florida. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. 1st edition. San Diego, CA: Academic Press, pp. 245–265.
- Shane S.H.** (1990b) Comparison of bottlenose dolphin behavior in Texas and Florida, with a critique of methods for studying dolphin behavior. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. 1st edition, San Diego, CA: Academic Press, pp. 541–558.
- Smolker R.A., Richards A.F., Connor R.C. and Pepper J.W.** (1992) Sex differences in patterns of association among indian ocean bottlenose dolphins. *Behaviour* 123, 38–69.
- Sutherland W.J.** (1998) The importance of behavioral studies in conservation biology. *Animal Behaviour*, 56, 801–809.
- Teixeira A.M. and Duguy R.** (1981) Observations de Delphinidés dans les eaux cotières portugaises. *Relatórios de Actividades do Aquário Vasco da Gama* 9, 1–9.
- Wells R.S. and Scott M.D.** (1994) Bottlenose dolphin. In Ridgway S.R. and Harrison F.R.S. (eds) *Handbook of marine mammals*. 2nd edition. London: Academic Press, pp. 137–182.
- Whitehead H.** (1995) Investigating structure and temporal scale in social organizations using identified individuals. *Behavioral Ecology* 6, 199–208.
- Whitehead H.** (1999) Testing association patterns of social animals. *Animal Behaviour* 57, 26–29.
- Whitehead H.** (2007) Selection of models of lagged identification rates and lagged association rates using AIC and QAIC. *Communications in Statistics: Simulation and Computation* 36, 1233–1246.
- Whitehead H.** (2008a) *Analyzing animal societies: quantitative methods for vertebrate social analysis*. 1st edition. Chicago: University of Chicago Press.
- Whitehead H.** (2008b) Precision and power in the analysis of social structure using associations. *Animal Behaviour* 75, 1093–1099.
- Whitehead H.** (2009) SOCPROG programs: analyzing social structures. *Behavioral Ecology and Sociobiology* 63, 765–778.
- Whitehead H. and Connor R.** (2005) Alliances I. How large should alliances be? *Animal Behaviour* 69, 117–126.
- Whitehead H., Bejder L. and Ottensmeyer C.A.** (2005) Testing association patterns: issues arising and extensions. *Animal Behaviour* 69, e1–e6.
- Würsig B. and Würsig M.** (1977) The photographic determination of group size, composition, and stability of coastal porpoises (*Tursiops truncatus*). *Science* 198, 755–756.
- and
- Zar J.H.** (1996) *Biostatistical analysis*. 4th edition. Upper Saddle River, NJ: Prentice-Hall.

#### Correspondence should be addressed to:

J.F. Augusto  
 Unidade de Investigação em Eco-Etologia  
 ISPA—Instituto Universitário  
 Rua Jardim do Tabaco 34, 1149-041 Lisboa, Portugal  
 email: jaugusto@ispa.pt



