

## **Nemo through the looking-glass: a commentary on Desjardins & Fernald**

Rui F. Oliveira and Adelino V. M. Canário

*Biol. Lett.* 2011 **7**, doi: 10.1098/rsbl.2010.0760 first published online 27 April 2011

---

### **References**

This article cites 10 articles, 2 of which can be accessed free  
<http://rsbl.royalsocietypublishing.org/content/7/4/487.full.html#ref-list-1>

### **Subject collections**

Articles on similar topics can be found in the following collections

[behaviour](#) (634 articles)  
[cognition](#) (126 articles)  
[neuroscience](#) (78 articles)

### **Email alerting service**

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

## Comment

**Nemo through the looking-glass: a commentary on Desjardins & Fernald**

Mirror-elicited aggression has been widely used in animal behaviour research. Recently, Desjardins & Fernald [1] reported that males of the cichlid fish *Astatotilapia burtoni* fighting a mirror had higher expression of immediate early genes in brain areas homologous to the amygdala and the hippocampus than males fighting a real opponent, despite the fact that the behavioural and androgen responses did not differ between the two conditions. Surprisingly, Desjardins & Fernald [1] omitted two recent papers (Oliveira *et al.* [2] in another cichlid fish, *Oreochromis mossambicus*, and Hirschenhauser *et al.* [3] in quail, *Coturnix japonica*), which also found a dissociation between the behavioural and the physiological responses towards mirror images. In these two studies, the mirror image elicited an aggressive response but failed to trigger the androgen response present when fighting a real opponent, suggesting that the physiological response to the fight depends on the cognitive appraisal that the individuals make of the situation [4]. This omission is even more surprising as the two divergent results support contrasting adaptive roles for socially driven changes in androgen levels (aka ‘challenge hypothesis’, [5]): the lack of hormonal response to a mirror fight suggests the need for asymmetric feedback to adjust androgen production, which is therefore only functional for subsequent interactions; the presence of an androgen response triggered by a mirror fight suggests a role for androgens in the modulation of aggressive behaviour in the ongoing fight. Both the *O. mossambicus* and quail studies support the first hypothesis, while the *A. burtoni* study is the first evidence for the latter.

Three different kinds of factors may contribute to the divergent androgen responses.

- *Phylogenetic bias.* Phylogeny is an unlikely source of variation in this case since two of the species showing divergent responses are closely related cichlid fishes, one of which shares the same pattern of response with the phylogenetically relatively distant quail.
- *Differences in the behavioural tasks.* In the real opponent treatment, Desjardins & Fernald [1] staged fights between males previously placed in separate compartments. Therefore, the fish interacting across the transparent partition were mimicking a territorial border fight between two territorial males, with no initial asymmetry in resource-holding potential and no clear outcome. In contrast, in the other two studies (Hirschenhauser *et al.* [6], was used as

a real opponent control for the mirror fights in Oliveira *et al.* [2]) a resident/intruder paradigm was used. This could explain why Desjardins & Fernald [1] did not find differences both in aggressive behaviour and in androgen levels between mirror and real fights, but it does not explain why mirror fights elicited a significant androgen response when compared with control males who had not been in an aggressive encounter.

- *Differences in hormone assays.* Androgen levels reported by Desjardins & Fernald [1] are much lower (ca. 2000 times) than those published for other cichlids, including those previously reported for the same species by the same laboratory [7]. Since Desjardins & Fernald [1] appear to have used blood plasma directly in the assay, and not the extracts of fluids (as done in the other studies, i.e. Oliveira *et al.* [2], Parikh *et al.* [7] and Hirschenhauser *et al.* [3]), the presence of androgen-binding globulins could have the effect of reducing apparent levels of androgens in unpredictable and not necessarily proportional ways. The biological meaning of changes at such a low level (less than 150 pg ml<sup>-1</sup>) is questionable, but assuming a hypothetical proportional distribution of the putative error among the three experimental groups of Desjardins & Fernald [1], the divergent results cannot be explained.

The fact that Desjardins & Fernald [1] assayed hormones from plasma, and in the other studies hormones were assayed from excreted metabolites in urine [2], or from droppings [3], raises the possibility that the delayed time for the excretion of hormones could be generating these differences. However, the sampling time used in both non-invasive studies has taken into account the known excretion time for these metabolites [3,8], and should reflect the circulating androgen levels at the end of the fights. Moreover, both studies did detect a response in the presence of real opponents using the same time course, and in cichlid fish, a significant positive correlation was detected between plasma and urine levels [9].

Finally, it should be noted here that in the target paper, the expression of the immediate early gene *egr-1* in the pre-optic area (POA) exhibited a non-significant trend (owing to high intra-treatment variation) to be higher in fish fighting real opponents than their mirror image. Since the expression of *egr-1* in response to social stimuli is highly co-localized with GnRH-1 neurons in the POA ( $R = 0.93$ , see Burmeister *et al.* [10]), these data suggest a higher activity of GnRH-1 neurons in fish fighting real opponents than in mirror fighters, which should translate into higher androgen levels in the former ones. Therefore, the *egr-1* expression data in the POA is in agreement with androgen levels reported in previous studies and contrasts with the ones reported in the target paper.

In summary, the divergent results between Desjardins & Fernald [1] and previously published studies could have a biological or artefactual basis and should not be ignored. The standardization of behavioural paradigms and hormone assay methods and

the integration of brain activation measures in future studies need to be considered in order to promote the integration of data and the establishment of general principles in the field, namely the adaptive function of socially driven changes in androgen levels.

The authors thank Katharina Hirschenhauser, Ton Groothuis, Peter Dijkstra, David Gonçalves, Magda Teles and Teresa Fagundes for discussing this topic with us. During the writing of this manuscript research undertaken by R.F.O. was supported by FCT (PTDC/PSI/71811/2006).

Rui F. Oliveira<sup>1,2,\*</sup> and Adelino V. M. Canário<sup>3</sup>

<sup>1</sup>*Instituto Superior de Psicologia Aplicada, Rua Jardim do Tabaco 34, 1149-041 Lisboa, Portugal*

<sup>2</sup>*Champalimaud Neuroscience Programme, Instituto Gulbenkian de Ciência, Rua da Quinta Grande 6, 2780-156 Oeiras, Portugal*

<sup>3</sup>*Centro de Ciências do Mar, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal*

\*Author for correspondence ([ruiol@ispa.pt](mailto:ruiol@ispa.pt)).

- 1 Desjardins, J. K. & Fernald, R. D. 2010 What do fish make of mirror images? *Biol. Lett.* **6**, 744–747. (doi:10.1098/rsbl.2010.0247)
- 2 Oliveira, R. F., Carneiro, L. A. & Canario, A. V. M. 2005 No hormonal response in tied fights. *Nature* **437**, 207–208. (doi:10.1038/437207a)
- 3 Hirschenhauser, K., Wittek, M., Johnston, P. & Möstl, E. 2008 Social context rather than behavioral output or winning modulates post-conflict testosterone responses in Japanese quail (*Coturnix japonica*). *Physiol. Behav.* **95**, 457–463. (doi:10.1016/j.physbeh.2008.07.013)
- 4 Oliveira, R. F. 2009 Social behavior in context: hormonal modulation of behavioural plasticity and social competence. *Integr. Comp. Biol.* **49**, 423–440. (doi:10.1093/icb/icp055)
- 5 Wingfield, J. C., Hegner, R. E., Dufty Jr, A. M. & Ball, G. F. 1990 The ‘challenge hypothesis’: theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *Am. Nat.* **136**, 829–846. (doi:10.1086/285134)
- 6 Hirschenhauser, K., Taborsky, M., Oliveira, T., Canario, A. V. M. & Oliveira, R. F. 2004 A test of the ‘challenge hypothesis’ in cichlid fish: simulated partner and territory intruder experiments. *Anim. Behav.* **68**, 741–750. (doi:10.1016/j.anbehav.2003.12.015)
- 7 Parikh, V. N., Clement, T. S. & Fernald, R. D. 2006 Androgen level and male social status in the African cichlid, *Astatotilapia burtoni*. *Behav. Brain Res.* **166**, 291–295. (doi:10.1016/j.bbr.2005.07.011)
- 8 Vermeirssen, E. L. M. & Scott, A. P. 1996 Excretion of free and conjugated steroids in rainbow trout (*Oncorhynchus mykiss*): evidence for branchial excretion of the maturation-inducing steroid, 17,20  $\beta$ -dihydroxy-4-pregnen-3-one. *Gen. Comp. Endocrinol.* **101**, 180–194. (doi:10.1006/gcen.1996.0020)
- 9 Hirschenhauser, K., Canario, A. V. M., Ros, A. F. H., Taborsky, M. & Oliveira, R. F. 2008 Social context may affect urinary excretion of 11-ketotestosterone in African cichlids. *Behaviour* **145**, 1367–1388. (doi:10.1163/156853908785765935)
- 10 Burmeister, S. S., Jarvis, E. D. & Fernald, R. D. 2005 Rapid behavioral and genomic responses to social opportunity. *PLoS Biol.* **3**, e363. (doi:10.1371/journal.pbio.0030363)